

## Review Article

# Organo-mineral associations in temperate soils: Integrating biology, mineralogy, and organic matter chemistry§

Ingrid Kögel-Knabner<sup>1\*</sup>, Georg Guggenberger<sup>2</sup>, Markus Kleber<sup>3</sup>, Ellen Kandeler<sup>4</sup>, Karsten Kalbitz<sup>5</sup>, Stefan Scheu<sup>6</sup>, Karin Eusterhues<sup>1,7</sup>, and Peter Leinweber<sup>8</sup>

<sup>1</sup> Lehrstuhl für Bodenkunde, Department für Ökologie und Ökosystemmanagement, Wissenschaftszentrum Weihenstephan, Technische Universität München, 85350 Freising-Weihenstephan, Germany

<sup>2</sup> Institut für Agrar- und Ernährungswissenschaften, Martin-Luther-Universität Halle-Wittenberg, Weidenplan 14, 06108 Halle/Saale, Germany

<sup>3</sup> Department of Crop and Soil Science, Oregon State University, 3017 Ag & Life Sciences Bldg., Corvallis, OR 97331–7306, USA

<sup>4</sup> Institut für Bodenkunde und Standortslehre, Fachgebiet Bodenbiologie, Universität Hohenheim, Emil-Wolff-Straße 27, 70599 Stuttgart, Germany

<sup>5</sup> Lehrstuhl für Bodenökologie, Universität Bayreuth, 95440 Bayreuth, Germany

<sup>6</sup> Institut für Zoologie, Technische Universität Darmstadt, Schnittspahnstraße 3, 64287 Darmstadt, Germany

<sup>7</sup> present address: Chemisch-Geowissenschaftliche Fakultät, Friedrich-Schiller-Universität, Burgweg 11, 07749 Jena, Germany

<sup>8</sup> Institut für Landnutzung, Universität Rostock, Justus-von-Liebig-Weg 6, 18051 Rostock, Germany

## Abstract

We summarize progress with respect to (1) different approaches to isolate, extract, and quantify organo-mineral compounds from soils, (2) types of mineral surfaces and associated interactions, (3) the distribution and function of soil biota at organo-mineral surfaces, (4) the distribution and content of organo-mineral associations, and (5) the factors controlling the turnover of organic matter (OM) in organo-mineral associations from temperate soils. Physical fractionation achieves a rough separation between plant residues and mineral-associated OM, which makes density or particle-size fractionation a useful pretreatment for further differentiation of functional fractions. A part of the OM in organo-mineral associations resists different chemical treatments, but the data obtained cannot readily be compared among each other, and more research is necessary on the processes underlying resistance to treatments for certain OM components. Studies using physical-fractionation procedures followed by soil-microbiological analyses revealed that organo-mineral associations spatially isolate C sources from soil biota, making quantity and quality of OM in microhabitats an important factor controlling community composition. The distribution and activity of soil microorganisms at organo-mineral surfaces can additionally be modified by faunal activities. Composition of OM in organo-mineral associations is highly variable, with loamy soils having generally a higher contribution of polysaccharides, whereas mineral-associated OM in sandy soils is often more aliphatic. Though highly reactive towards Fe oxide surfaces, lignin and phenolic components are usually depleted in organo-mineral associations. Charred OM associated with the mineral surface contributes to a higher aromaticity in heavy fractions. The relative proportion of OC bound in organo-mineral fractions increases with soil depth. Likewise does the strength of the bonding. Organic molecules sorbed to the mineral surfaces or precipitated by Al are effectively stabilized, indicated by reduced susceptibility towards oxidative attack, higher thermal stability, and lower bioavailability. At higher surface loading, organic C is much better bioavailable, also indicated by little <sup>14</sup>C age. In the subsurface horizons of the soils investigated in this study, Fe oxides seem to be the most important sorbents, whereas phyllosilicate surfaces may be comparatively more important in topsoils. Specific surface area of soil minerals is not always a good predictor for C-stabilization potentials because surface coverage is discontinuous. Recalcitrance and accessibility/aggregation seem to determine the turnover dynamics in fast and intermediate cycling OM pools, but for long-term OC preservation the interactions with mineral surfaces, and especially with Fe oxide surfaces, are a major control in all soils investigated here.

**Key words:** OM / organic matter stabilization / microhabitat / fractionation / specific surface area / NMR / SEM / <sup>14</sup>C age

Accepted July 13, 2007

\* Correspondence: Prof. Dr. I. Kögel-Knabner;  
e-mail: koegel@wzw.tum.de

§ **Topical Issue** *Soils as a source and sink for CO<sub>2</sub> – Mechanisms and regulation of organic matter stabilisation in soils* (editors: I. Kögel-Knabner and E. Matzner). Synthesis of the DFG Priority Program SPP 1090 (German Research Foundation—“Deutsche Forschungsgemeinschaft”).



## 1 Introduction

Inverse correlations between grain size and organic carbon (OC) content have frequently been ascribed to stabilization of organic matter (OM) in soils by close associations between OM and mineral surfaces (Tieszen et al., 1984; Anderson, 1988; Oades, 1988), giving rise to the paradigm of sorptive protection. Mineral surfaces in soils are mainly provided by the very small particles (<2 µm) that constitute the soil clay fraction. This fraction is a mixture of phyllosilicates, oxides, and hydroxides, and, in some soil types, short-range order minerals like allophane and imogolite. Traditional concepts were highly focused on the role of phyllosilicates as dominating mineral surfaces for the formation of organo-mineral associations (e.g., Stevenson, 1982; Theng, 1979). Later, total mineral surface area has been considered as a predictor for the amount of OM that is stabilized in soils (Mayer et al., 1994a; Saggart et al., 1996; Hassink, 1997; Six et al., 2002), but an understanding on the relationship between mineralogy and the chemistry of OM bound in organo-mineral associations is lacking. Most of the available data are from loamy arable topsoils, and information on other soil types and the distribution with depth still is lacking (von Lützow et al., 2006).

A major step forward was achieved from the combined efforts of soil-mineralogical, organic-chemical, and microbial research approaches. Increasing evidence emerged on the important influence of soil biota in the formation of organo-mineral assemblages (Chenu and Stotzky, 2002). Here, the ability of microorganisms to standardize mineral surfaces to their needs by “active support preconditioning” (Bos et al., 1999; Dufrene et al., 1996) seems to promote mineral-organic associations through the deposition of extracellular polysaccharides and proteins as adhesives. It is well-known that numerous Gram-negative and Gram-positive bacteria exude extracellular polymeric substances (EPS) into their environment (Omoike and Chorover, 2006). In natural aqueous environments, pristine mineral surfaces become coated rapidly by biogenic organic films (Bos et al., 1999). The formation of these “conditioning films” moderates eventual differences in the surface chemistry of the underlying substrate and can thus be seen as an adaptation mechanism that allows microorganisms to colonize highly variable types of mineral surfaces.

Spatial variability has historically been regarded as random noise in the system, but it becomes now evident that soil microorganisms as well as soil enzymes are heterogeneously distributed within the soil matrix (Ettema and Wardle, 2002; Kandeler and Dick, 2006; Fry, 2007). Still little detailed information is available on the spatial distribution of soil microbiota. This spatial variability and heterogeneity in the distribution of microorganisms and OM with regard to the mineral surfaces requires new approaches to the investigation of possible interactions between OM and mineral surfaces.

In this paper, we summarize the recent results obtained within the PP1090 “Soils as a source and sink for CO<sub>2</sub> – mechanisms and regulation of organic matter stabilisation in soils” and discuss our current understanding of sorptive organo-

mineral associations. *A priori* the topics discussed in this manuscript focus on mechanisms at the organic matter–mineral interface. Firstly, we approach this by studies on the separation/isolation or extraction of organo-mineral associations and the quantity of organo-mineral compounds in soils. Then we highlight recent research achievements with respect to the types of mineral surfaces and the associated interactions. Based on that, distribution and content of OM in different soils, its composition and stability in organo-mineral associations are discussed in light of the distribution of microorganisms and microbial enzyme activities at organo-mineral surfaces in soils. For a discussion of protection of OC due to occlusion of OM within clay microstructures and aggregates see Bachmann et al. (2008, this issue, pp. 14–26) and Flessa et al. (2008, this issue, pp. 36–51). A detailed description of the soils and sites investigated in the PP1090 is given by Kögel-Knabner et al. (2008, this issue, pp. 5–13).

## 2 Approaches to isolate, extract, and quantify organo-mineral compounds in soils

### 2.1 Physical separation methods

Mineral-bound OM has been defined as all OM that is adsorbed to minerals or entrapped in small micro-aggregates (Wattel-Koekkoek et al., 2004; Chenu and Plante, 2006). To make quantitative estimates of the amount and stability of OM in organo-mineral associations *sensu strictu* and to investigate both their mineral and their organic composition, it is necessary to specifically isolate this fraction from soils. This has been approached by physical aggregate disruption in combination with particle-size and/or density separation or by chemical treatments that are assumed to specifically either attack or isolate organo-mineral components.

Density fractionations have been used for many years in attempts to separate free OM (mostly plant residues) from OM bound to minerals. Organo-mineral fractions obtained from the physical-fractionation approach are generally isolated after disruption of aggregates into so-called primary particles (Christensen, 1992, 1996) and subsequent particle and density fractionation (von Lützow et al., 2006, 2007). Two techniques are used for aggregate disruption and dispersion, sonication at different energies (Amelung and Zech, 1999) or end-over-end agitation in water with glass beads (Balesdent et al., 1991). North (1976) suggested to check complete dispersion by calibration of a clay-sized fraction obtained in proportion to that achieved during standard particle-size analyses. The different aggregate stabilities and densities of organo-mineral associations in soils of different mineralogy have to be considered, when adjusting a fractionation procedure to a specific soil (Turchenek and Oades, 1979; Schmidt et al., 1999a, b). This explains why there is until now no accepted standard protocol with respect to, e.g., input energy to disrupt the aggregates or the critical densities of the heavy liquid to separate the light fraction from mineral-associated OM. Chenu and Plante (2006) showed that many of the so-called <2 µm “particles” were in fact nanometer- to micrometer-sized micro-aggregates in which OM was

encrusted by minerals or coated minerals. None-the-less, young plant residues are often well separated from older and thus stable organo-mineral associations by a combination of ultrasonic dispersion followed by particle-size and density fractionation (Turchenek and Oades, 1979; Golchin et al., 1994; Wander, 2004; Schöning and Kögel-Knabner, 2006; Kaiser and Guggenberger, 2007b). Some authors do not use any dispersion procedure prior to density fractionation (Solins et al., 2006; Janzen et al., 1992). Although this avoids the artificial inclusion of more mineral material in the light fraction (Janzen et al., 1992), it does not allow a differentiation in primary and secondary particles in the sense of Christensen (1992, 1996).

## 2.2 Chemical extraction and degradation methods

Various extraction methods using desorbing, hydrolyzing, and oxidizing reagents have been used to extract or enrich OM fractions associated with the soil mineral phase in the framework of the PP1090 (Ellerbrock et al., 2005; Eusterhues et al., 2003, 2005a; Kleber et al., 2005; Mikutta et al., 2006; Helfrich et al., 2007). For a detailed survey of existing procedures see von Lützow et al. (2007).

Organic matter extracted from soil with alkaline Na-pyrophosphate (pH 9–10) is supposed to be liberated from metal-organic associations and from mineral surfaces *via* desorption. But Na-pyrophosphate not only liberates metals from metal-organic complexes but also extracts polymeric metal species and small-sized minerals associated with OM like goethite, ferrihydrite, amorphous Al(OH)<sub>3</sub>, gibbsite, and layered silicates (Schuppli et al., 1983; Kassim et al., 1984; Adams and Kassim, 1984; Evans and Wilson, 1985; Kaiser and Zech, 1996). Hence, the suitability of pyrophosphate extraction to reliably recover stable OM from soils still remains to be demonstrated, especially in comparison with other more established methods like hydrolysis or oxidation treatments.

Acid hydrolysis using hot 6 N HCl is a common procedure for the quantification of old, biochemically resistant OM fractions in soils (Paul et al., 1997, 2001; Huang et al., 1999; Collins et al., 2000). This is mainly due to the preferential removal of labile O-alkyl C, primarily originating from polysaccharides, and the selective enrichment of more resistant aromatic and aliphatic C components (Dieckow et al., 2005). However, fresh plant detritus needs to be removed prior to hydrolysis because plant residues partly resist hydrolysis, which could bias the assessment of stable C in soils. But even then acid hydrolysis does not unambiguously isolate a fully stable OM pool (Balesdent, 1996; Plante et al., 2006). One should further bear in mind that at low pH, Fe oxides are dissolved and, thus, OM stabilized *via* interaction with this part of the mineral phase may get released.

Chemical degradation of soil OM by oxidizing reagents is an alternative method to preferentially remove OM structures unprotected by the mineral matrix. But it oxidizes preferentially chemically labile components such as polysaccharides and aromatics (Leifeld and Kögel-Knabner, 2001; von Lützow et al., 2007). The organic residuum has been shown to be

composed of compounds inherently stable against chemical attack and protected by association with soil minerals (reviewed by Mikutta et al., 2005). For example, fossil or black C (Schmidt et al., 1999c) and aliphatic compounds such as n-alkanes and n-fatty acids likely originating from aliphatic biopolymers (Griffith and Schnitzer, 1977; Martin et al., 1981; Schulten et al., 1996; Cuypers et al., 2002) comprise oxidation-resistant OM (Eusterhues et al., 2005a). The concentration of oxidation-resistant OM in various soils can vary strongly from 1% to 80% with usually larger relative amounts in subsoils than in topsoils (Eusterhues et al., 2003, 2005a; Kaiser and Guggenberger, 2003). The authors assigned this increasing resistance of OM to oxidation with soil depth to a stronger binding to the mineral phase at smaller surface loading of the minerals by OM. At present, it is still a drawback of the oxidation procedures that they are hardly comparable due to differences in methodology, *i.e.*, oxidant reagent concentration, temperature, solid-to-solution ratios, and number of repetitions, and due to the fact that certain OM components may be removed in preference to others (*e.g.*, lignin-derived aromatics; Mikutta et al., 2006). Future research needs to focus on the comparability of methods and on the understanding of underlying processes that cause certain OM components to resist extraction, hydrolysis, or oxidative degradation.

## 3 Types of mineral surfaces and associated interactions

Three types of functional surface sites on minerals can be distinguished, which offer different bonding possibilities to the OM. Their availability is assumed to control area density as well as the strength of the bonds formed: (1) Surfaces inhabited by single coordinated hydroxyl groups are common to Fe and Al oxides, allophane, and imogolite and the edges of layer silicates; (2) siloxane surfaces with permanent layer charge due to isomorphous substitution. In soils, these surfaces appear at some 1:2-layer-type phyllosilicates, such as vermiculite, illite, and smectite; (3) siloxane surfaces without layer charge. This surface type occurs in both 1:2-layer-type (talc, pyrophyllite) and in 1:1-type minerals (kaolinite group). The results from the PP1090 largely follow this energetic scheme: Strong protection was generally correlated with the presence of hydroxylated mineral phases in acidic soil environments.

In bulk soils and clay fractions of illitic, near-neutral soils, correlations between SOC and CEC (Kahle et al., 2002, 2003a) were described. In accordance with the above theoretical considerations, these findings were interpreted to be caused by cation exchange as the dominating bonding mechanism between mineral matrix and OM. The fact that no or negative correlations exist between the Fe of total Fe oxides (Fe<sub>o</sub>) and TOC was explained by the presence of well-crystalline and therefore less reactive Fe oxides in the investigated soils or by the decreased number of positively charged surface hydroxyls in near neutral environments. But there is evidence for a specific interaction between Fe oxides and OM in soils of similar mineralogy and pH (Luvisol data, Fig. 7 and 8; Schöning et al., 2005b). Most often information on the binding mechanisms is obtained from theoretical considerations, pro-

cesses found in short-term laboratory experiments, or from statistical correlations. Further studies are needed to directly investigate the specific binding mechanisms found in organo-mineral associations isolated from soils using high-resolution spectroscopic techniques.

Oxidation-resistant OM is a chemically stable, old OM fraction (Theng et al., 1986). In a wide range of different soils, its concentration often correlates with the content of Fe oxides and/or clay (Eusterhues et al., 2003) or Fe oxides and short-range order Al silicates (Kleber et al., 2005; Mikutta et al., 2006). In soils with low amounts of poorly crystalline components ( $Fe_o : Fe_d < 0.2$ ), the mineral-associated stable OC relates positively to the content of crystalline Fe oxides (Mikutta et al., 2006; Fig. 1). These results strongly suggest that in acidic soil environments, stable OM is predominantly associated with Fe oxides and short-range order Al silicates. Such minerals provide large reactive surface areas (Eusterhues et al., 2005b), which renders them most suitable for interaction with OM either via sportive interactions or coprecipitation. In addition to strong bonds due to ligand exchange, an inherent recalcitrance of the oxidation-resistant OM fraction may lead to its oxidation resistance.

As preferential sorption sites the mouth of micropores (<2 nm) or small mesopores (2–10 nm) of Fe oxides have been suggested (Kaiser and Guggenberger, 2003). Besides of the boundaries of individual domains, these small pores represent the surface roughness resulting from defects in the surface structure of goethite crystallites (Weidler et al., 1998). Consequently, micropores are sites of increased reactivity

and thus best suited for sorptive interactions with organic acids and polyelectrolytes (Kaiser and Guggenberger, 2003; Mikutta et al., 2004). There, OM can form strong multiple bindings that disfavor both the desorbability and the oxidative removal of OM, respectively (Kaiser and Guggenberger, 2007a). Similarly, Zimmerman et al. (2004) observed larger desorption hystereses of nitrogenous compounds following sorption to mesoporous sorbents compared with their nonporous analogues. However, in soil and sediment environments, the importance of molecular-sized pores within the mineral matrix for the stabilization of OM is not yet understood. Mayer et al. (2004) concluded that other factors than association with mesopores must be responsible for the accumulation of OM in marine sediments because only <10%–20% of OM was found to be actually contained in 2–10 nm pores that are small enough to prevent access of hydrolytic enzymes and microbes. Similar results are reported by Kaiser and Guggenberger (2007a) for the soil mineral goethite. Close to sorption maximum, the volume ratio between goethite and OM is almost 1. At this surface loading, the configuration of OM is less dense than at lower loading, and OM bridges goethite crystals and crystallites to form small aggregates (Kaiser and Guggenberger, 2007a). This type of OM with less organic ligands being involved in the bonding is prone to desorption and oxidative degradation, and possibly also to a better biological decomposition.


## 4 Distribution of microorganisms and microbial enzyme activities at organo-mineral surfaces in soils

### 4.1 Methods used to study small-scale distribution of soil microorganisms

Different methods have been applied to study the distribution of OM and microorganisms in the soil matrix, such as electron microscopy, repeated washing of soil aggregates, and techniques based on physical soil fractionation (Stemmer et al., 1998). Using biological soil thin sections and a combination of image analysis and geostatistical tools, the presence of spatial patterns in the distribution of bacteria was demonstrated at the microscale, with ranges of spatial autocorrelation of 1 mm and below (Nunan et al., 2003). For detailed microvisualization, optical-microscope techniques are applied, supported by specific labeling techniques and image analysis (cf., review by Schmid et al., 2006; Watt et al., 2006). The confocal laser-scanning microscope is a very useful tool to document the spatial arrangement of microbes and their *in situ* activities, as indicated by fluorescence-labeling and reporter technique. Physical fractionation of soils offers the opportunity to study compartmentation of substrates and soil microorganisms at the small scale (Kandeler et al., 2005; Marx et al., 2005).


### 4.2 Fungi are more important/abundant in coarse fractions

In general, organic particles >2000  $\mu\text{m}$ , which consist predominantly of particulate residues from plant material, decompose very rapidly while the OM in smaller-size fractions is




**Figure 1:** Three-dimensional plot showing the relation of the content of mineral-protected organic C (MOC) with the contents of poorly crystalline minerals and metal-humic complexes  $[(Fe + Al)_o]$  and crystalline Fe oxides ( $Fe_{d-o}$ ). In samples 1–5, where little poorly crystalline minerals are present ( $Fe_o : Fe_d < 0.2$ ), a significant relationship of MOC with  $Fe_{d-o}$  ( $r^2 = 0.93$ ) suggests that MOC is mainly associated with crystalline Fe oxides. Arrows are displayed for better visualization only (taken from Mikutta et al., 2006).

characterized by an increase of the degree of degradation and a decrease in the C : N ratio. The dominance of fungal-derived xylanase activity in coarse fractions has recently been documented for both bulk soil and earthworm casts (Fig. 2, from Marhan et al., 2007). Xylanase activity and fungal-derived phospholipid fatty acids show a very close relationship to the particulate OM of the coarse-sand fraction (Kandeler et al., 1999a, b, c, 2000, 2001; Stemmer et al., 1999; Poll et al., 2003). The close vicinity of fungi, extracellular xylanase, and particulate OM guarantees an efficient use of the substrates by the fungal consumer (Kandeler and Dick, 2006). These results were also supported by the highly significant correlation between xylanase activity and the quality of OM (the signal of hemicelluloses measured by pyrolysis mass spectrometry (Fig. 3)).



**Figure 2:** Distribution of xylanase activity in particle-size fractions of a Terra-fusca Rendzina from a 130 y-old beech forest near Göttingen (southern Lower Saxony, Germany). Addition of beech leaf litter (44.5% C, 1.6% N) induced mainly an increase of xylanase activity in the sand fractions (from Marhan et al., 2006).



**Figure 3:** The response of xylanase activity of a Haplic Phaeoem to the abundance of its corresponding substrate. Substrate quality of hemicellulose was measured by its xylan-to-xylose ratio calculated from indicator signals of Py-FIMS. Soil samples (0–20 cm) from the control (unfertilized) and NPK plots of the “Eternal Rye Cultivation” were incubated for 2, 7, 21, and 35 d at 25°C. Xylanase activity was expressed as µg glucose equivalents (GE) per gram of soil and 24 h (Leinweber et al., unpublished).

### 4.3 Bacteria dominate the small-size fractions

Small-sized fractions contain the most microbial biomass in different soils (Kanazawa and Filip, 1986; Jocteur Monrozier et al., 1991). For example, cellulose-feeding bacteria (*Cytophaga*, *Cellulomonas*) attach closely with their substrate in these microhabitat, which minimizes the loss of enzymes into the soil solution by diffusion (Alexander, 1977). Therefore, enzymes involved in the hydrolysis of high molecular weight C substrates reflect the location and perhaps the quality of OM in agricultural soils (Speir and Ross, 2002). The bacterial dominance in the finer fraction was also shown by high abundance of bacterial-derived phospholipid fatty acids, high diversity of 16s rRNA gene fragments, and high activities of enzymes involved in the cycling of N and P (Kandeler et al., 2000; Poll et al., 2003). Clay-sized particles have a higher surface area than coarser particles, which facilitates bacterial growth as well as attachment and protection of microorganisms and extracellular enzymes. Using molecular techniques (RFLP—restriction fragment length polymorphism), we found, that the silt- and clay-sized fraction favored a greater richness of bacterial species (Sessitsch et al., 2001; Kirchmann et al., 2004; Selesi et al., 2007). PLFA pattern of particle-size fractions support these results (Poll et al., 2003) showing a rather simple fungal-dominated community in sand-sized fractions and complex bacterial-dominated communities in silt- and clay-sized fractions. Since <0.01% of the soil surface is colonized, spatial isolation of organisms might promote richness of microbial diversity in these fractions. Further evidence for greater habitat diversity in small-sized soil fractions comes from studies of Sessitsch et al. (2001) who found aerobic and strictly anaerobic bacteria species in the clay-sized fraction and no strictly anaerobic species in sand-sized fractions.

The high affinity of bacteria to small particle size fractions, as described above, suggests that mineral-organic associations in these fractions show a “microbial fingerprint”. Indications for this are low C : N ratios as commonly observed for bacterial biomass (Knicker, 2004), enrichment in <sup>13</sup>C (Gleixner et al., 1993), and a microbial signature of the polysaccharides in the fine fractions (Guggenberger et al., 1994; Kiem and Kögel-Knabner, 2003).

Although studies using particle-size fractions helped to understand the location and functioning of soil microorganisms at the small scale, the fractionation procedure destroys the natural micro-environment of particles and soil microorganisms (especially branching of hyphal organisms) and does not provide any insight into soil microbial processes in the three-dimensional soil system (Kandeler and Dick, 2006). This disadvantage can partly be overcome by applying slicing techniques of soil cores followed by microanalytical methods and modeling (Gaillard et al., 2003; Poll et al., 2006). First attempts with these approaches are experimental designs focusing on the description and modeling of two-dimensional gradients of microbial activities that occur in the detritusphere as well as in preferential-flow paths of soils (Gaston and Locke, 2002; Poll et al., 2006, 2007). In a model experiment, bacteria and fungi showed differing utilization strategies of organic substrates at the soil-litter interface. Coupling PLFA


analysis with stable isotope techniques, *Poll et al.* (2006) revealed that fungi assimilated C directly in the litter, whereas bacteria took up the substrates in the soil and therefore depended more on transport processes than fungi (*cf.*, also *Beare et al.*, 1997; *Ekschmitt et al.*, 2008, this issue, pp. 27–35).

#### 4.4 Soil animals as drivers of the small-scale distribution of soil microorganisms and microbial-enzyme activities at organo-mineral surfaces

By mixing organic and mineral soil components during the gut passage, decomposer soil invertebrates affect the distribution of microorganisms among soil particle-size fractions, their activity, and, therefore, mobilization and stabilization processes of OM (*Scullion and Malik*, 2000; *Marhan and Scheu*, 2006). A large number of soil invertebrates are involved in these processes but due to their large size and dominance in numbers and biomass in many terrestrial ecosystems, earthworms are the most important drivers (*Beare et al.*, 1995; *Bardgett*, 2005). Both sorption and desorption of OM from mineral surfaces occurs during the gut passage through earthworms (*Shipitalo and Protz*, 1988, 1989; *Hindell et al.*, 1997; *Marhan and Scheu*, 2006). By ingesting and processing large amounts of mineral soil, endogeic earthworms in particular affect mobilization and stabilization processes of OM (*Kubierna*, 1948; *Shaw and Pawluk*, 1986; *Scheu*, 1987). By the addition of large amounts of mucus and water in the foregut, aggregates are broken up and clay minerals are dispersed, thereby liberating physically protected OM and making it accessible for digestion in the mid-gut of the animal (*Barois*, 1992; *Schmidt et al.*, 1999d). On the contrary, by resorption of water and mucus in the hindgut, earthworms stimulate the re-association and new formation of organo-mineral complexes thereby contributing to the stabilization of OM. Desorption and resorption processes during the gut passage result in the frequently observed pattern of

increased decomposition of OM in fresh cast materials but reduced decomposition in aging casts (*Martin*, 1991; *Lavelle et al.*, 2004; *Tiunov and Scheu*, 2000; *Marhan and Scheu*, 2006).

Although the gut passage through earthworms strongly affects soil structure, microbial activity, and decomposition processes, the composition of the microbial community appears to be little affected. Fluorescence microscopy, genetic fingerprinting (T-RFLP analysis), and phospholipid fatty analysis suggests that the general structure of microbial communities remains little affected by the gut passage through earthworms (*Schönholzer et al.*, 1999; *Egert et al.*, 2004; *Marhan et al.*, 2007). However, by feeding on both OM-rich patches and mineral soil and intimately mixing them during the gut passage, the structure and activity of microbial communities are strongly affected (“nutrient-enrichment effects”; *Devliegher and Vertraete*, 1995). Further, despite the structure of microbial communities may be little affected, their activity may change markedly, *e.g.*, it has been documented recently that the activity of denitrifiers is strongly increased in earthworm guts and fresh cast materials (*Horn et al.*, 2006; *Drake and Horn*, 2006). Further, although little information is available, the few studies undertaken suggest that the structure of microbial communities associated with particle-size fractions in soil is little modified by the gut passage through earthworms (*Marhan et al.*, 2007; *Sampedro et al.*, 2006; Fig. 4). Rather than changing the associations of microorganisms with certain particle-size fractions, earthworms alter the distribution of microorganisms associated with soil macro- and micro-aggregates (*Mummey et al.*, 2006), and this likely is responsible for modifying the decomposition of OM enclosed in earthworm casts in the long term (*Tiunov and Scheu*, 2000; *Marhan and Scheu*, 2006). The formation of micro-aggregates within macro-aggregates and an associated microbial community in the micro-aggregates which is relatively inactive compared to macro-aggregate populations likely contributes to the long-term stabilization of OM in earthworm casts (*Mummey et al.*, 2006).



**Figure 4:** Xylanase activity in bulk soil and particle-size fractions of soil and casts of *Lumbricus terrestris* without (– Litter) and with addition of beech litter (+ Litter). Means of two or three replicates  $\pm 1$  S.D. GE: glucose equivalents (from *Marhan et al.*, 2006).

## 5 Amount and distribution of mineral-bound OM in soils

The amount and relative percentage of OC that is found in organo-mineral fractions is highly variable between soil types, but also between horizons within the same soil. Only a limited number of studies exist that give a complete data set for the OC concentration as well as the distribution of OC between fractions. Table 1 gives a summary from studies with available data from a number of different soils using the dispersion and density fractionation approach. The relative proportion of the OC that is associated with the heavy fraction is highly variable. Organic C concentrations in the heavy fraction vary between  $<1$  and  $>100$  g OC kg<sup>-1</sup>. This may be due to a number of reasons. Organic C distribution might be affected by different procedures for aggregate disruption as mentioned above. Likewise, different densities of 1.6 to 2.2 g cm<sup>-3</sup> for separation of the clay-bound OC are used. As pointed out by Christensen (1992), Vanyushina and Travnikova (2003), and Sollins et al. (2006), this strongly affects the mass ratio of organic to mineral phase of these particles. The differences in proportion of OC in the heavy fractions may be due to variations in mineralogy (amount and type of minerals) and OC loading of the minerals (Oades, 1988; Christensen, 1996; Hedges and Oades, 1997). So, a soil richer in short-range-ordered Fe oxyhydroxides and Al silicates shows larger concentrations in mineral-bound fractions than that richer in clay minerals (Rasmussen et al., 2005). In the comparison of different density separates obtained without dispersion, Sollins et al. (2006) showed that primary minerals with a low affinity to OM predominate in the fractions  $>2.28$  g cm<sup>-3</sup>. Carbon-14 data show that the OC associated with these seemingly unreactive minerals has longer turnover times than the OC in the lighter fractions which are supposed to include the more reactive mineral surfaces.


Generally, the relative proportion of OC bound in organo-mineral fractions is lower in forest topsoils compared to soils under agriculture (e.g., John et al., 2005; Eusterhues et al., 2005a, b). In agricultural soils, the lower input of OM and the frequent disruption of aggregates (Six et al., 1998) leads to a preferential loss of OC from free and occluded particulate OM fractions. This is seen in the data for the Rottalmünster site with different land use in Tab. 1. Similar data with a preferential loss of free OC in the clay-sized fraction are described by Chenu and Plante (2006). Table 1 also shows that the relative importance of clay-bound OC increases with soil depth. It becomes especially high in subsoil horizons with high contents of clay-sized minerals, such as Bt or spodic horizons. In the subsoil horizons, in particular different OC loading of the mineral phase may be responsible for the different OC concentrations in the heavy fractions, in addition to the variable mineralogy (Tab. 1). But only limited data are available for the OC loading of the mineral phase, i.e., the ratio of OC to the specific surface area of the minerals.

## 6 Specific surface area of the mineral phase and loading with OM

Particle-size fractionation of the PP1090 soils exhibits highest C concentrations in the fine fractions  $<6.3$   $\mu\text{m}$  (Rumpel et al.,

2004). This observation matches the widely accepted assumption that minerals in the clay fraction stimulate C storage and protection from degradation. It is also in accord with the idea that the available inorganic surface area controls OM sorption. Organic C loadings in the range of 0.5 to 1 mg OC m<sup>-2</sup> were reported in sediments and soil (Keil et al., 1994; Mayer, 1994b and references therein) and suggested to represent a monolayer coverage over all mineral surfaces. Later, Mayer and Xing (2001) found the extent of surface loading in acid soils to depend on pH, with higher loadings (close to 1 mg OC m<sup>-2</sup>) at low pH, and interpreted this as evidence for a patchy distribution of OM on mineral surfaces. However, detailed investigations of the silt- and clay-sized minerals in the PP1090 soils led us to different conclusions:

Kahle et al. (2003b) and Kleber et al. (2004) found higher Fe oxide densities in coarse-clay versus fine-clay fractions to correspond with higher OC loading in coarse- versus fine-clay fractions. This indicates the ability of minerals to preserve OC results from the combined influence of reactive surface sites and a large specific surface area. Figure 5 shows the OC loading for a number of soils investigated in the course of the SPP differentiated into (1) A horizons from acidic forest soils (Waldstein, Steinkreuz, Borsberg, Zeughaus), (2) in B and C horizons of these same soils, and (3) in A horizons from circumneutral, mainly illitic arable soils (Halle, Rottalmünster). It shows a decline of OC loadings with depth, obviously mirroring OC contents and thus suggesting a strong coupling of OC loadings to OM input (Kaiser and Guggenberger, 2003). Only the A horizons have a loading higher than the so-called monolayer equivalent. Scanning



**Figure 5:** Organic matter loadings for fine-clay and coarse-clay particle-size fractions. Error bars represent mean values and the standard error (SE) of the means. The dotted horizontal line indicates the loading level commonly taken to represent a monolayer equivalent of organic matter over all surfaces and was termed the monolayer-equivalent (ME) level (Mayer, 1994a). The term monolayer-equivalent is stressed, because it is reasonably well established that organic matter covers mineral surfaces in a discontinuous manner (Ransom et al. 1998; Arnarson and Keil, 2001; Mayer and Xing, 2001; Kahle et al., 2002).

**Table 1:** Amount and distribution of light and heavy OM fractions in different soils.


Reference	Comments	Soil type	Description	Depth or horizon	fPOM (C in fraction % of SOC)	oPOM (C in fraction % of SOC)	Mineral (C in fraction % of SOC)	Free (g kg <sup>-1</sup> )	Occluded (g kg <sup>-1</sup> )	Mineral (g kg <sup>-1</sup> )
<i>Roscoe and Buurman</i> (2003)	D = 1.7 g cm <sup>-3</sup>	Typic Haplustox	Cerrado	0–7.5 cm	25.9	40.5	33.7	25.8	40.4	33.6
		Typic Haplustox	after conversion into cropland, no tillage	0–7.5 cm	23.9	45.8	30.3	21.2	40.7	26.9
		Typic Haplustox	after conversion into cropland, plow tillage	0–7.5 cm	25.1	45.5	29.4	24.0	43.6	28.2
		Typic Haplustox	Cerrado	7.5–15 cm	29.8	41.6	28.6	28.3	39.5	27.1
		Typic Haplustox	after conversion into cropland, no tillage	7.5–15 cm	29.4	42.7	27.8	27.3	39.6	25.8
		Typic Haplustox	after conversion into cropland, plow tillage	7.5–15 cm	26.7	43.5	29.8	25.2	41.1	28.2
<i>Rasmussen et al.</i> (2005), <sup>14</sup> C analyses	D = 1.6 g cm <sup>-3</sup>	Ultic Haploxeralfs	Granite soil	A2	37.6	24.7	37.6	14.0	9.2	14.0
		Ultic Haploxeralfs	Granite soil	Bt	5.4	6.3	88.2	4.8	5.6	78.0
		Ultic Haploxeralfs	Granite soil	BC	18.8	47.8	33.3	1.3	3.3	2.3
		Ultic Haploxeralfs	Andesite-Granite soil	A2	30.6	18.6	50.8	22.5	13.7	37.4
		Ultic Haploxeralfs	Andesite-Granite soil	Bt	17.0	24.4	58.7	4.8	6.9	16.6
Ultic Haploxeralfs	Andesite-Granite soil	BC	17.9	50.0	32.1	3.0	8.4	5.4		
<i>John et al.</i> (2005)	D = 2.0 g cm <sup>-3</sup> , oPOM originally determined for two density fractions	Luvisol (Rottalmünster)	Wheat	0–30 cm	3.3	10.1	86.6	0.4	1.2	10.3
		Luvisol (Rottalmünster)	Maize	0–30 cm	4.1	9.1	86.8	0.5	1.2	11.2
		Luvisol (Rottalmünster)	Grass	0–10 cm	3.7	10.3	85.9	0.9	2.6	21.5
		Luvisol (Rottalmünster)	Grass	10–20 cm	3.6	5.4	91.0	0.4	0.6	10.6
		Luvisol (Rottalmünster)	Grass	20–30 cm	5.9	6.2	87.9	0.5	0.5	7.5
		Luvisol (Rottalmünster)	Forest	0–7 cm	33.6	18.0	48.4	13.5	7.3	19.5
		Luvisol (Rottalmünster)	Forest	7–25 cm	20.4	16.8	62.8	1.9	1.6	6.0
		Luvisol (Rottalmünster)	Forest	25–40 cm	14.6	8.2	77.2	0.5	0.3	2.5
<i>Swanston et al.</i> (2005)	D = 1.6 g cm <sup>-3</sup> , <sup>14</sup> C analyses	Ultisol	Forest	0–15 cm	25.5	16.7	49.7			
		Ultisol	Forest	15–30 cm	23.8	14.9	51.9			
<i>Echeverria et al.</i> (2004)	D = 2 g cm <sup>-3</sup>	Hapludults, Kanhapludults	Eatonton	0–10 cm or A horizon	10.2		89.8			
		Kanhapludults	Athens	0–10 cm or A horizon	4.8		95.2			
		Paleudults, Haplohumods	Waycross	0–10 cm or A horizon	12.1		87.9			
		Haplothords	N. Florida	0–10 cm or A horizon	14.7		85.3			
<i>Rovira and Vallejo</i> (2003)			mediterranean calcareous forest soils	H	89.2	4.7	6.1			

Reference	Comments	Soil type	Description	Depth or horizon	fPOM (C in fraction % of SOC)	oPOM (C in fraction % of SOC)	Mineral (C in fraction % of SOC)	Free (g kg <sup>-1</sup> )	Occluded (g kg <sup>-1</sup> )	Mineral (g kg <sup>-1</sup> )
				mediterranean calcareous soils	A	57.8	16.8	25.4		
				mediterranean calcareous soils	B	26.1	17.2	56.8		
Golchin et al. (1994)	D = 1.6 g cm <sup>-3</sup> , NMR	Solodic (Natrixeralf)	mixed forest with grass ground cover	0–10 cm	19.3	9.2	71.5			
		Noncalic brown earth	(Hapludalf)	Poplar box forest with blue and wallaby grasses	0–10 cm	14.7	16.6	68.7		
			Grey Clay (Chromustert)	Brigalow forest	0–10 cm	31.3	17.5	51.2		
			Red-brown earth (Rhodoxeralf)	Old pasture (>60 yrs)	0–10 cm	7.7	9.7	82.6		
			Black earth (Pellustert)	Native grassland	0–10 cm	6.9	15.6	77.5		
Eusterhues and Kögel-Knabner (unpublished)	D = 2 g cm <sup>-3</sup>	Cambisol (Steinkreuz)	acid forest under beech	0–5 cm or Ah horizon			10.0			8.3
		Cambisol (Steinkreuz)	acid forest under beech	5–24 cm or Bv horizon			38.0			3.7
		Cambisol (Steinkreuz)	acid forest under beech	24–50 cm or SdBv1 horizon			42.0			3.4
		Cambisol (Steinkreuz)	acid forest under beech	50–80 cm or SdBv2 horizon			68.0			2.0
		Cambisol (Steinkreuz)	acid forest under beech	85–115 cm or IIICv horizon			89.0			1.0
		Cambisol (Steinkreuz)	acid forest under beech	115–140 cm or IVCv1 horizon			73.0			0.8
Eusterhues and Kögel-Knabner (unpublished)	D = 2 g cm <sup>-3</sup>	Podzol (Waldstein)	acid forest under spruce	0–6 cm or Aeh horizon			19.0			7.2
		Podzol (Waldstein)	acid forest under spruce	6–9 cm or Bsh horizon			27.0			25.1
		Podzol (Waldstein)	acid forest under spruce	9–10 cm or Bs horizon			33.0			17.2
		Podzol (Waldstein)	acid forest under spruce	10–47 cm or Bv1 horizon			98.0			7.8
		Podzol (Waldstein)	acid forest under spruce	47–60 cm or Bv2 horizon			75.0			1.5
		Podzol (Waldstein)	acid forest under spruce	60–100 cm or Cv horizon			73.0			1.5
Kaiser and Guggenberger (unpublished)	D = 1.6 g cm <sup>-3</sup>	Cambisol (Steinkreuz)	acid forest under beech	0–5 cm or Ah horizon	65.0	12.0	23.0	45.3	8.4	16.0
		Cambisol (Steinkreuz)	acid forest under beech	5–24 cm or Bw1 horizon	40.0	10.0	50.0	4.4	1.1	5.5
		Cambisol (Steinkreuz)	acid forest under beech	24–50 cm or Bw2 horizon	15.0	8.0	77.0	0.5	0.3	2.7
		Cambisol (Steinkreuz)	acid forest under beech	50–80 cm or Bw3 horizon	10.0	9.0	81.0	0.2	0.1	1.3
		Cambisol (Steinkreuz)	acid forest under beech	80–85 cm or 2C horizon	5.0	7.0	88.0	0.1	0.1	1.3
		Cambisol (Steinkreuz)	acid forest under beech	85–115 cm or 3C horizon	10.0	6.0	84.0	0.1	0.1	1.2


Reference	Comments	Soil type	Description	Depth or horizon	fPOM (C in fraction % of SOC)	oPOM (C in fraction % of SOC)	Mineral (C in fraction % of SOC)	Free (g kg <sup>-1</sup> )	Occluded (g kg <sup>-1</sup> )	Mineral (g kg <sup>-1</sup> )
		Cambisol (Steinkreuz)	acid forest under beech	115–140 cm or 4C horizon	10.0	5.0	85.0	0.1	0.0	0.7
	D = 1.6 g cm <sup>-3</sup>	Podzol (Waldstein)	acid forest under spruce	0–10 cm or Aeh horizon	67.0	8.0	25.0	24.6	3.0	9.5
		Podzol (Waldstein)	acid forest under spruce	10–12 cm or Bh horizon	16.0	11.0	73.0	15.0	10.3	68.1
		Podzol (Waldstein)	acid forest under spruce	12–30 cm or Bs horizon	9.0	12.0	79.0	4.7	6.2	41.0
		Podzol (Waldstein)	acid forest under spruce	30–55 cm or Bw horizon	7.0	5.0	88.0	0.5	0.4	6.7
		Podzol (Waldstein)	acid forest under spruce	55–70 cm or C1 horizon	5.0	5.0	90.0	0.1	0.1	2.1
		Podzol (Waldstein)	acid forest under spruce	70–80 cm or C2 horizon	5.0	3.0	92.0	0.1	0.0	1.4
<i>Baisden et al. (2002)</i>	fPOM oPOM < 1.6–2.2 g cm <sup>-3</sup> DF > 2.22 g cm <sup>-3</sup>	Sierra Nevada	annual grassland	0–2.5 cm	45.0	40.0	8.0	26.6	23.6	4.7
				2.5–9 cm	27.0	54.0	17.0	7.8	15.7	4.9
				9–22 cm	3.0	49.0	57.0	0.4	5.9	6.8
		Sierra Nevada	annual grassland	0–2 cm	27.0	34.0	20.0	4.1	5.2	3.1
				2–10 cm	10.0	35.0	43.0	0.8	2.6	3.2
				10–29 cm	7.0	35.0	51.0	0.2	0.9	1.3
		Sierra Nevada	annual grassland	0–2 cm	54.0	30.0	6.0	10.8	6.0	1.2
				2–12 cm	11.0	50.0	24.0	6.2	28.0	13.4
				12–24 cm	7.0	27.0	56.0	0.1	0.5	1.1
		Sierra Nevada	annual grassland	0–2 cm	21.0	43.0	26.0	12.0	24.5	14.8
				2–11 cm	9.0	24.0	66.0	0.9	2.3	6.3
				11–40 cm	2.0	9.0	88.0	0.1	0.4	3.9
<i>Chenu and Plante (2006)</i>	tPOM 1.6 g cm <sup>-3</sup> organo-mineral: >1.6 g cm <sup>-3</sup> and <2.2 g cm <sup>-3</sup>	France	forest under pine	0–30 cm	2.3		86–83	66.0		104.0
		France	maize since 35 years, converted from forest	0–30 cm	1.4		84–76	16.0		41–42

electron microscopy of the fine fractions shows that the OM is not uniformly distributed over all clay-mineral surfaces: whereas most phyllosilicate surfaces are free of OM, some platelets are continuously coated across their basal surface (Fig. 6, Eusterhues et al., unpublished, *cf.*, also *Bachmann et al., 2008*, this issue, pp. 14–26). Accordingly, we could not find strong correlations between the mineral-bound organic C

and the total specific mineral surface (Fig. 7a, *Eusterhues et al., 2005b*). On the other hand, strong correlations exist between the mineral-bound OC and the Fe oxide content for the subsoils (Fig. 7b; *Eusterhues et al., 2005b*; unpublished). This suggests that Fe oxides are the dominant carrier for mineral-bound OM in the subsoils. In contrast to subsoil horizons, A horizons of all soils contain too much C or insufficient




**Figure 6:** SEM micrograph of silt and clay particles from the topsoil of the Rottalmünster Luvisol: (a) secondary electron image; (b) EDX dot map of C distribution of image a.



**Figure 7:** Relation between mineral-bound OM and (a) specific surface area or (b) Fe oxide content.  $OC_{\text{mineral}}$  is the organic-C concentration of the density fraction  $>2 \text{ g cm}^{-3}$ ,  $SSA_{\text{minerals}}$  the BET- $N_2$  surface area measured after destruction of the organic matter with  $Na_2S_2O_8$ ,  $Fe_{\text{DCB}}$  is the concentration of dithionite-citrate-bicarbonate-extractable Fe.

Fe oxides to follow the observed trend. In the topsoil, we either assume a larger C loading on Fe oxides or a greater importance of other mineral surfaces. As can be deduced from the sorption experiments of OM on goethite and ferrihydrite, the capacity of these minerals to protect sorbed OM is smaller than their capacity to bind it (Kaiser and Guggenberger, 2007a; Kaiser et al., 2007). Different C-to-FeDCB ratios between topsoils and subsoils or between the acid forest subsoils and the neutral arable subsoil can be caused by differences in (1) the sorbate composition, (2) the amount of protonated hydroxyl groups at different pH, (3) the OM input, and (4) the specific Fe oxide-surface area (Kaiser and Guggenberger, 2003; Eusterhues et al., 2005b; Kleber et al., 2005). Mass balance calculations, trying to reconstruct the measured total mineral-surface area by summing up assumed specific surface areas of the individual mineral phases (cf., Eusterhues et al., 2005b for details) indeed yield varying specific surface areas for ferrihydrite (300–800  $\text{m}^2 \text{g}^{-1}$ ) and goethite (25–300  $\text{m}^2 \text{g}^{-1}$ ) in different soils. In the Luvisol, the lowest specific surface areas of 300  $\text{m}^2 \text{g}^{-1}$  for ferrihydrite and 25  $\text{m}^2 \text{g}^{-1}$  for goethite were sufficient to reconstruct the measured specific surface area. In Fig. 8, the total mineral-bound OC is related to these calculated surface areas of ferrihydrite and goethite. Regression lines for the subsoils of all soils show similar slopes, quantifying the OM loading within the range of 1.1 to 1.9 mg C per  $\text{m}^2$  oxide surface. These loadings must be regarded as maximum values since a certain fraction is additionally fixed on the phyllosilicates and probably also on poorly crystalline allophane-like phases. However, these loadings are higher but in the same magnitude than compared to maximum loadings found for sorption of the so-called hydrophilic fraction of DOM on goethite (0.22–0.50  $\text{mg C m}^{-2}$  goethite) and the so-called hydrophobic fraction of DOM (1.13–1.28  $\text{mg C m}^{-2}$  goethite; Kaiser and Guggenberger, 2003). The mono-layer-equivalent coating is assumed to be approx. 1  $\text{mg C m}^{-2}$  (Keil et al., 1994; Mayer, 1994a), while Bergamaschi et al. (1997) found typical loadings of 2–5  $\text{mg m}^{-2}$  for sediments underlying high-productivity zones. Murphy et al. (1990) report a Langmuir-type isotherm for the sorption of peat humic acid on hematite, showing its sorption maximum at 1.1  $\text{mg C m}^{-2}$  hematite. However, at high concentrations of humic acid (approx. 5–20  $\mu\text{mol mL}^{-1}$ ), the sorption plateau passes into a rising slope again, corre-



**Figure 8:** Relation between mineral-bound OM and specific surface area of Fe oxides.


sponding to, e.g., 2.8 mg C m<sup>-2</sup> hematite at 20 μmol mL<sup>-1</sup>. This was interpreted as reflecting multiple layering of humic acids on the mineral surface.

The fact that all subsoils, though different in Fe oxide content, pH, and pedogenesis, show similar C loadings suggests that 1–2 mg C m<sup>-2</sup> might be a typical OC loading for Fe oxides in soil. We concluded that Fe oxides are the most important sorbents for the formation of organo-mineral associations in the subsoil, while phyllosilicate surfaces are comparatively less important.

## 7 Composition of OM in organo-mineral associations

Methods that can be applied to solid samples, such as analytical pyrolysis, (thermo)chemolysis, and <sup>13</sup>C-NMR spectroscopy are particularly useful to characterize the composition of the OM associated with the minerals. Generally differences of bulk-soil OM from different soils are not very pronounced (Mahieu et al., 1999), and bulk soil OM does also not reflect

the different plant input. To identify the contribution of plant litter input, the particulate OM fractions are most useful. Figure 9 for the fractions from Rotthalmünster shows a plant debris composition, dominated by high proportions of O-alkyl C (mainly cellulose and hemicelluloses), followed by alkyl C, and smaller contributions of aromatic C, resembling the composition of the respective site-specific plant residues. Compared to that, the mineral-bound OM is generally depleted in lignin and phenolic components (Guggenberger et al., 1994; Kiem and Kögel-Knabner, 2003; Schöning et al., 2005a). This is also seen in the <sup>13</sup>C-NMR spectra for all soils investigated here (Fig. 9), as the signal intensity in the aromatic region (110–160 ppm) and especially the aryl-O signal at 150 ppm characteristic for phenolic compounds is low. Baldock et al. (1992) and Mahieu et al. (1999) already pointed out that clay-size fractions generally show a higher content of alkyl C than the whole soils and in particular the coarser fractions, as revealed by <sup>13</sup>C-NMR spectroscopy. Generally the fine or heavy fractions show higher proportions of carboxyl C than plant residues, indicative of the more oxidized stage of the stabilized OM. The fine fractions isolated from loamy soils are dominated by high proportions of O/N alkyl C, as demon-



**Figure 9:** Composition of light and heavy fractions of organic matter isolated from the sites Bad Lauchstädt, Waldstein, Steinkreuz, and Rotthalmünster as obtained from the analysis by <sup>13</sup>C-NMR spectroscopy (from Kiem et al., 2003; Rumpel et al., 2004; Helfrich et al., 2006).




Figure 9: continued




Figure 9: continued





Figure 9: continued

strated for the Rotthalmünster soils. They are mostly assigned to polysaccharides and proteins, with large contributions of microbial polysaccharides (Guggenberger et al., 1994; Kiem and Kögel-Knabner, 2003). In contrast, the OM in the fine fractions of sandy soils are dominated by long-chain aliphatic molecules in the topsoil horizons, illustrated for both acid sandy soils Waldstein and Steinkreuz by the signal at 30 ppm. At present, it is not clear, what the mechanism underlying these differences in the composition of sandy and loamy soils is. In the subsoil, the fine fraction of the sandy acid forest soils is composed of low molecular weight organic acids, as indicated by the dominance of alkyl C in combination with the high carboxylic signal intensity around 175 ppm in the  $^{13}\text{C}$ -NMR spectra. This reflects the formation of the acid subsoil horizons via a chromatographic process that accumulates small, highly oxidized organic acids in the deeper horizons (Rumpel et al., 2004).

In soils containing charred OM, a part of this char is found in association with the mineral phase, as can be delineated from  $\text{HNO}_3$  oxidation and SEM (Brodowski et al, 2005, 2007).



**Figure 10:** Thermograms of bulk samples (a) and of phenols and lignin monomers (b) of dissolved organic matter (DOM) from the Oi and Oa horizon of a podzolic soil and from maize straw before and after sorption to a mineral soil. Data for DOM before sorption are derived from Kalbitz et al. (2003) whereas data after sorption are unpublished. These data were obtained in a similar way like in Kalbitz et al. (2003). The mean residence times (MRT) of the bulk samples were published by Kalbitz et al. (2005).

This leads to higher proportions of aromatic C with a high signal intensity around 130 ppm in the  $^{13}\text{C}$ -NMR spectra as evident in the  $^{13}\text{C}$ -NMR spectra from Bad Lauchstädt (Fig. 9), while other analytical techniques, such as pyrolysis or thermochemolysis, are often not able to detect the charred OM.

## 8 Stability of OM in organo-mineral associations

### 8.1 Stabilizing effect of binding of OM to mineral surfaces: evidence from sorption experiments

Sorption to minerals generally reduces the susceptibility of OM towards oxidative attack (Kaiser and Guggenberger, 2003) and the bioavailability (Kalbitz et al., 2005). Interactions between soil minerals and DOM result in a distinct stabilization as indicated by increased mean residence times after sorption to a mineral subsoil (Kalbitz et al., 2005), soil minerals (Mikutta et al., 2007), and after precipitation by Al (Scheel et al., 2007) (Fig. 10a). So, the mineralization rate constants of a stable C pool ( $k_2$  of a double-exponential model) increased by a factor between 3 and 25 comparing DOM and sorbed OM (Fig. 10a; Kalbitz et al., 2005). This increase in stability against microbial decay was accompanied by an increase in thermal stability as shown by pyrolysis–field ionization mass spectrometry (Py-FIMS). The major volatilization peak of bulk DOM shifted by 200 K, 255 K, and 125 K after sorption of DOM from maize straw, Oi and Oa horizons, respectively. The volatilization curves of main compound classes (carbohydrates, phenols and lignin monomers, lipids, alkylaromatics, heterocyclic N-containing compounds, peptides) showed a thermal behavior similar to the bulk DOM (Fig. 10b). This indicates that inherently labile organic compounds of DOM such as carbohydrates are stabilized by soil minerals to a similar extent as more stable components like phenols and lignin monomers. Kalbitz et al. (2005) found that labile DOM from maize straw or Oi horizons was stabilized by soil minerals to a larger extent than already stable DOM from Oa horizons. This observation coincides with a larger shift in the Py-FIMS volatilization peak after sorption of this labile

DOM (maize, Oi: +200 to +255 K) in comparison to sorption of stable DOM (Oa: +125 K). That means, the extent of stabilization against microbial decomposition was indicated by the extent in the shift towards higher temperatures after sorption. This is in line with *Buurman et al. (2002)* who explained high oxidation temperatures of OM from strongly developed profiles with high C-to-Fe and C-to-sesquioxide ratios by the stabilization of OM in OM-metal precipitates and in differences in organic chemistry. Coincidence of thermal stability and resistance against microbial decomposition was also reported in other studies by Py-FIMS (*Schulten and Leinweber, 1999, 2000*). This is explained by a correlation between the activation energy required for the thermal bond cleavage in Py-FIMS and the chemical energy required for the enzymatic cleavage of chemical bonds in microbial-decomposition processes. From these results, we can conclude that stabilization of DOM by organo-mineral interactions can be traced by two independent methods, measurements of C mineralization in incubation experiments and analysis of the thermal behav-

ior using Py-FIMS. More studies are required to get deeper insight into the energetic details of OM bonds, mineral associations, and microbial decomposition. *Mikutta et al. (2007)* showed that OM bound to minerals mainly by ligand exchange was more resistant against mineralization than OM held by noncoulombic interactions (van der Waals forces). Furthermore, Ca bridges enhanced the stability of sorbed OM, but less than the binding *via* ligand exchange (*Mikutta et al., 2007*).

### 8.2 Stabilizing effect of binding of OM to mineral surfaces: age and turnover of OM in organo-mineral associations isolated from soils

When comparing the <sup>14</sup>C ages of the density fractions <1.6 g cm<sup>-3</sup> with those of the fractions >1.6 g cm<sup>-3</sup> (Tab. 2), it is obvious that the <sup>14</sup>C age of the latter is generally higher, with the exception of the three uppermost horizons at Wald-

**Table 2:** Organic carbon concentrations and storage in different fractions of the soils Steinkreuz and Waldstein, along with the <sup>14</sup>C activity in percent modern carbon and <sup>14</sup>C age in years before 1950 (*Eusterhues et al., 2003, 2005; Kaiser and Guggenberger, unpublished*); numbers in brackets indicate OC percentage in the fractions related to bulk soil OC.

Horizon	Depth	OC concentration	OC storage	<sup>14</sup> C activity bulk soil	<sup>14</sup> C age bulk soil	<sup>14</sup> C activity of H <sub>2</sub> O <sub>2</sub> -resistant fraction	<sup>14</sup> C age of H <sub>2</sub> O <sub>2</sub> -resistant fraction	<sup>14</sup> C activity of fraction oxidized by H <sub>2</sub> O <sub>2</sub> <sup>a</sup>	<sup>14</sup> C age of fraction oxidized by H <sub>2</sub> O <sub>2</sub> <sup>a</sup>	<sup>14</sup> C activity of <1.6 g cm <sup>-3</sup>	<sup>14</sup> C age of <1.6 g cm <sup>-3</sup>	<sup>14</sup> C activity of >1.6 g cm <sup>-3</sup>	<sup>14</sup> C age of >1.6 g cm <sup>-3</sup>	<sup>14</sup> C activity of fraction >1.6 g cm <sup>-3</sup> and resistant to NaOCl treatment <sup>a</sup>	<sup>14</sup> C age of fraction >1.6 g cm <sup>-3</sup> and resistant to NaOCl treatment <sup>a</sup>
	(cm)	(g kg <sup>-1</sup> )	(kg m <sup>-2</sup> )	(pm C)	(y BP)	(pm C)	(y BP)	(pm C)	(y BP)	(pm C)	(y BP)	(pm C)	(y BP)	(pm C)	(y BP)
Steinkreuz (Cambisol)															
A	0–5	82.6	4.29	112.3	modern	104.5	modern	112.9	modern	112.4	modern	111.9	modern	99.3	60±30 (1)
Bw1	5–24	9.8	2.41	101.3	modern	86.6	1150±22 (5)	101.8	modern	104.8	modern	98.0	160±25 (50)	72.3	2606±35 (5)
Bw2	24–50	3.0	0.79	92.1	655±25	71.8	2665±35 (16)	95.9	340 (84)	119.0	modern	84.2	1375±30 (77)	36.2	8170±80 (12)
Bw3	50–80	1.4	0.40	80.9	1700±30	51.5	5325±45 (27)	92.0	672 (73)	122.8	modern	69.8	2890±30 (81)	31.0	9400±240 (24)
3C	85–115	1.1	0.14	80.6	1758±56	49.3	5675±45 (30)	93.1	571 (70)	117.2	modern	70.8	2780±45 (84)	21.6	12310±170 (31)
4C1	115–140	0.5	0.07	76.3	2165±30	66.8	3245±40 (58)	88.7	959 (42)	116.4	modern	69.1	2960±30 (85)	21.4	12390±190 (34)
Waldstein (Podzol)															
EA	0–10	38.1	2.90	93.6	525±30	88.1	1015±25 (3)	94.2	479 (97)	92.0	655 (75)	98.5	120±25 (25)	90.7	835±25 (3)
Bh	10–12	92.8	1.04	98.5	120±25	91.4	725±20 (6)	99.5	40 (94)	95.8	435 (27)	99.5	30±20 (73)	85.7	1290±25 (15)
Bs	12–30	52.0	5.47	91.1	745±40	72.6	2575±25 (9)	92.8	597 (91)	87.0	1010 (21)	92.2	700±25 (79)	81.1	1715±20 (24)
Bw	30–55	7.7	2.09	82.2	1570±25	73.1	2515±30 (13)	83.3	1471 (87)	87.3	980 (12)	81.5	1640±20 (88)	70.5	2805±35 (38)
C1	55–70	1.7	0.25	62.0	3840±70	55.7	4700±35 (25)	64.1	3577 (75)	90.8	730 (10)	58.8	4265±30 (90)	43.7	6650±90 (47)
C2	70–80	1.9	0.19	62.0	3840±70	52.0	5260±45 (25)	65.4	3412 (75)	112.8	modern	56.5	4580±30 (92)	42.3	6910±110 (52)

<sup>a</sup> calculated by difference

stein. *Rasmussen et al.* (2005) also reported higher  $^{14}\text{C}$  ages for mineral-associated OM than for free particulate OM. Investigating six density fractions from  $<1.65\text{ g cm}^{-3}$  to  $>2.55\text{ g cm}^{-3}$ , *Sollins et al.* (2006) found too much charcoal in the two lightest fractions to allow for a reliable turnover estimate, but a continuous increase in turnover time from 150 to  $>980\text{ y}$  with increasing density in the subsequent density fractions. The relatively low  $^{14}\text{C}$  activity in the two lightest density fractions is probably caused by charcoal from a forest fire (*Sollins et al.*, 2006). Black C particles were frequently observed in the light fraction from the Waldstein surface-soil horizons and likely contribute to the relatively high  $^{14}\text{C}$  age in this fraction (*Marschner et al.*, 2008, this issue, pp. 91–110). In the PP1090 forest soils, the proportion of total OC and  $^{14}\text{C}$  age within the heavy fraction ( $>1.6\text{ g cm}^{-3}$ ) increased, concurrent with the increase of the  $^{14}\text{C}$  age of the oxidation-resistant fraction with soil depth. Such results were reported by a number of studies (e.g., *Torn et al.*, 1997; *Paul et al.*, 2001; *Swanston et al.*, 2005) and are indicative of increased mineral association and protection of OM at greater depths.

As noted above, extraction of OM provides no unambiguous results due to its unspecific nature. Nevertheless, combined information from various extracts together with information of the mineral assemblage may provide some useful information on the nature of the organic-mineral association. Sodium pyrophosphate has been applied sequentially after preliminary extraction of soil with water (*Ellerbrock et al.*, 2005) and NaOH (*Wattel-Koekkoek and Buurman*, 2004) in order to obtain more stable OM fractions. For instance, the pyrophosphate-soluble OM from two agricultural soils (Rotthalmünster and Halle) has been found to exhibit a slower turnover than the water-extractable OM (*Ellerbrock et al.*, 2005). Similarly, pyrophosphate-extractable OM from Mozambiquan clay fractions was found to contain older and probably more recalcitrant organic constituents than the OM extracted by NaOH (*Wattel-Koekkoek and Buurman*, 2004). Pyrophosphate has also been used to study the impact of metal-organic complexes on the stabilization of OM. For a set of terrace soils including Alfisols, Inceptisols, and Mollisols, *Masiello et al.* (2004) observed close correlations of the  $^{14}\text{C}$  inventories with both, pyrophosphate-extractable Al and Fe ( $r = 0.98$  and  $0.95$ , respectively). These correlations have been taken as evidence that complexation of OM by Al and Fe is an important mechanism for the stabilization of OM.

*Helfrich et al.* (2007) applied stepwise hydrolysis with trifluoroacetic acid to two agricultural soils (Rotthalmünster and Halle) after removal of particulate and humified OM and found also an enrichment of old OM in the sample residuum as reflected by  $^{13}\text{C}$  and  $^{14}\text{C}$  isotope patterns. The increase in the apparent  $^{14}\text{C}$  age of OM following hydrolysis was comparable to those observed after oxidative treatments using  $\text{H}_2\text{O}_2$ ,  $\text{Na}_2\text{S}_2\text{O}_8$ , and NaOCl ( $>1000\text{ y}$ ). These results fit quite well to earlier data from a Saskatchewan arable soil (Canada), showing that OM being resistant to  $6\text{ N HCl}$  hydrolysis is oldest and concurs with the  $^{14}\text{C}$  age of coarse clay and fine silt (*Anderson and Paul*, 1984). According to *Paul et al.* (2006), the nonhydrolyzable fraction shows an average of  $1,200\text{ y}$  greater mean residence times than total SOC.

Chemical degradation with  $\text{H}_2\text{O}_2$  (Tab. 2) was used in two acid sandy soils under forest to isolate a stable, residual OM fraction (*Eusterhues et al.*, 2003, 2005a). Generally, the  $^{14}\text{C}$  activity was smaller for the  $\text{H}_2\text{O}_2$ -resistant fraction than for the fraction oxidized by  $\text{H}_2\text{O}_2$ , meaning that in the resistant fraction, OC has a noticeably higher mean age. With increasing soil depth, the contribution of the  $\text{H}_2\text{O}_2$ -resistant fraction increased, as did the mean  $^{14}\text{C}$  age. This could indicate a more effective stabilization mechanism in the subsoil horizons.

Treatment of the  $>1.6\text{ g cm}^{-3}$  fraction with NaOCl left a fraction behind that is oldest amongst all fractions isolated by the different techniques (*Kaiser and Guggenberger*, unpublished; Tab. 2). The proportion of this old C fraction on total OC increased from  $<5\%$  in the surface soil to approx.  $50\%$  in the deepest horizons, again showing the pronounced protection of OM in the subsoil. However, a  $^{14}\text{C}$  age of  $>12,000\text{ y}$  in the NaOCl-resistant fraction of the 3C and 4C1 horizons at Steinkreuz may indicate the presence of geogenic OC in the Triassic sedimentary parent material. Such high  $^{14}\text{C}$  ages of NaOCl-resistant mineral-associated OM from subsoil horizons confirms the conclusion of *Kaiser and Guggenberger* (2003) that potential OM stabilization at mineral surfaces is limited to those OM molecules that sorb to surfaces with small surface loading. Turnover rates of OM are probably related to the time when the organic molecules are sorbed; the molecules that are sorbed first are best stabilized. Also *Mikutta et al.* (2006) reported  $^{14}\text{C}$  ages in a range of millenia for the NaOCl-resistant fraction of 12 bulk soils, while those of the NaOCl-removable fraction is mostly modern. Such differences indicate that mineral-associated OM may not just represent a stable, passive C pool but also a more active cycling one. This is corroborated by *Swanston et al.* (2005) who made use of an inadvertent  $^{14}\text{C}$ -pulse label to a whole forest system, caused by the incineration of medical waste. While in an uncontaminated control soil,  $^{14}\text{C}$  activity in the dense fraction was much less than in the free and occluded light fractions,  $^{14}\text{C}$  from the pulse label was rapidly incorporated into the dense fraction, showing that this organo-mineral fraction comprises highly stable material as well as more recent inputs (*Swanston et al.*, 2005). This is also obvious by mean residence time estimates for mineral-bound OM employing the  $^{13}\text{C}$ -natural abundance approach which is usually in the range of decades to a few centuries (*Skjemstad et al.*, 1990; *Balesdent et al.*, 1998). In contrast to the  $^{14}\text{C}$  method, the  $^{13}\text{C}$ -natural abundance approach gives an estimate of turnover dominated by relatively recent inputs and OC fractions that cycle within the time frame of the experiment (*Six and Jastrow*, 2002).

### 8.3 Stabilizing effect of binding of OM to surfaces: conceptual models accounting for varying degrees of sorptive preservation

The formation of stable OM with turnover times of up to several millenia by strong sorptive bondings to specific mineral surfaces is well-accepted, but the nature of the more rapidly cycling mineral-bound OM fraction is currently under discussion. According to *Kaiser and Guggenberger* (2003, 2007a),

at higher OM loadings at the mineral surface, less organic ligands per organic molecule are involved in the bonding, making the organic molecules more flexible and extending more into the soil solution, thus being more susceptible for microbial decomposition. *Ellerbrock et al. (2005)* assume that mainly the interaction between hydrophilic functional groups within OM and polyvalent cations located at mineral surfaces result in OM-mineral complexes, which may protect OM against decomposition. Therefore, the OM-to-clay ratio, the presence (or absence) of polyvalent cations, and the OM composition may influence distribution and conformation of OM on mineral surfaces. *Sollins et al. (2006)* and *Kleber et al. (2007)* provide an alternative explanation stating a zonal structure for OM associated with mineral surfaces, with the outer layers of the OM having a more rapid turnover than the inner ones. And finally, microbial colonization of mineral particles must be considered. *Swanston et al. (2005)* argue that to the extent that the heavy fraction acts as a sink for microbial biomass and products, it must include an OC component that can cycle rapidly.

## 9 Relevance of organo-mineral interactions for C stabilization in soils—conclusions and future research perspectives

We are still far away from agreed protocols for the quantitative/complete isolation or extraction of the mineral-bound OM from soils. Conventional physical fractionation protocols have to be tested for more soils and compared with the results from the newly emerging chemical approaches. Approaches combining physical-fractionation procedures with chemical treatments might allow for isolating mineral-bound fractions more specifically.

There is a clear improvement in our understanding of how soil microorganisms and their food sources are distributed within microhabitats. Clay-sized organo-mineral associations are preferred habitat for a complex microbial community offering substrates as well as high surface area for adsorption and protection of soil microorganisms. The richness of the soil microbial community in these fractions is due to the spatial isolation of organisms and substrates of different quality. Future studies should not only consider microbial-mediated decomposition of OM in different fractions, but also microbial-mediated changes of mineral surfaces which contribute to C stabilization of OM. Surface conditioning of mineral surfaces by microorganisms, which actively excrete polysaccharides and proteins warrants more attention for the creation of stable organo-mineral associations, as suggested by high contributions of microbial-derived organic compounds and low C : N ratios in the fine fractions. Since current studies in soil ecology focused mainly on soil fractions destroying the natural micro-environment of particles and soil microorganisms, a further challenge is to understand the distribution, structure, and function of soil biota in the three-dimensional soil system.

The proportion of OC bound to minerals and stored in the clay fraction increases with increasing depth and is generally higher in the subsoil compared to the topsoil. Likewise, the

strength of the bonding increases with soil depth, indicating the important role of subsoil OM in the OM stabilization. Clear differences are found between different soil types with respect to the type of OM in organo-mineral associations, although the underlying principles are not yet fully understood. Major components of the mineral-bound OM are of the alkyl and O/N-alkyl C type, whereas aromatic C is of minor importance, except for the presence of charred OM. In sandy acid soils, there is clear evidence for the stabilization of mainly alkyl C, which becomes more acidic and of lower molecular weight with increasing depth. Here, the stabilization seems to be mediated mainly *via* interaction with Fe oxides and short-range order Al silicates. In loamy soils, we find a much larger contribution of O/N alkyl C compounds, most probably microbial polysaccharides and proteins. This shows the importance of soil genesis on OM stabilization by formation of organo-mineral associations. At present, we have only a small data basis on the amount of OC in organo-mineral associations in selected soil types and horizons. A major task for future work is to enlarge our knowledge to more soil types, which also means a larger range in mineralogy.

Pedogenic oxides dominate the reactive surface area for organo-mineral interactions in all temperate soils investigated here. They seem to be especially important for the stabilization of OM in the subsoil. In arable topsoils developed from illitic loess, permanently charged siloxane surfaces of phyllosilicates are of additional relevance, although their specific surface area is generally much lower, compared to pedogenic Fe oxides. Total specific surface area, micropores, and mesopores could not be successfully used to predict OM accumulation by organo-mineral associations. This is most probably due to the fact that only a part of the mineral surfaces comes in contact with OM and that OM may not enter into small pore spaces. However, not the whole mineral-bound OM fraction shows the same (radiocarbon age, turnover, or) resistance against microbial degradation, so that we assume that effective stabilization only takes place for a certain fraction of the attached OM. It has also become clear that dissolved Al is a strong protective agent in acid soils. This mechanism is rather specific as restricted to acidic environments, but here, precipitation of OM by free Al is a significant control on OM preservation.

Major difficulties in the understanding and prediction of OM dynamics originate from the simultaneous operation of several mechanisms in addition to the protection by association of OM with mineral surfaces. This may be especially the case in A horizons, whereas we can often find the organo-mineral associations to be the dominating mechanism in subsoil horizons. Much more research is needed on the match between organic phase and mineral phase in relation to different soil types and conditions prevailing in these soils. Also, factors that control surface loading and coverage of minerals in different soil types need more attention in future work. We expect that identification and quantitation of the actual bonding mechanisms might explain a great deal of our data. For soil material this, however, is extremely challenging due to the heterogeneity and “imperfection” of the natural mineral surfaces as well as due to the small size of clays and oxides.

## Acknowledgments

We thank the “Deutsche Forschungsgemeinschaft” (DFG) for financial support of this work in the framework of the Priority Program 1090 “Soils as source and sink of CO<sub>2</sub> – mechanisms and regulation of organic matter stabilisation in soils”. We further would like to express our thanks for the possibility to work at several long-term experimental sites, which were provided by the Bayreuth Center of Ecology and Environmental Research, University of Bayreuth (BayCEER), the University of Halle-Wittenberg, The Helmholtz Centre for Environmental Research Leipzig-Halle (UFZ), and the “Höhere Landbauschule Rothalmünster”. We would also like to express our thanks to Margit von Lützwow and three anonymous reviewers for helpful comments and suggestions.

## References

- Adams, W. A., Kassim, J. K. (1984): Iron oxyhydroxides in soils developed from Lower Paleozoic sedimentary rocks in mid-Wales and implications for some pedogenetic processes. *J. Soil. Sci.* 35, 117–126.
- Alexander, M. (1977): Microbiology of the rhizosphere, in Alexander, M.: Introduction to Soil Microbiology. 2nd edn., John Wiley & Sons, New York, pp. 423–437.
- Amelung, W., Zech, W. (1999): Minimisation of organic matter disruption during particle-size fractionation of grassland epipedons. *Geoderma* 92, 73–85.
- Anderson, D. W. (1988): The effect of parent material and soil development on nutrient cycling in temperate ecosystems. *Biogeochem.* 5, 71–97.
- Anderson, D. W., Paul, E. A. (1984): Organo-mineral complexes and their study by radiocarbon dating. *Soil Sci. Soc. Am. J.* 48, 298–301.
- Arnarson, T. S., Keil, R. G. (2001): Organic-mineral interactions in marine sediments studied using density fractionation and X-ray photoelectron spectroscopy. *Org. Geochem.* 32, 1401–1415.
- Bachmann, J., Guggenberger, G., Baumgartl, T., Ellerbrock, R. H., Urbanek, E., Goebel, M.-O., Kaiser, K., Horn, R., Fischer, W. R. (2008): Physical carbon-sequestration mechanisms under special consideration of soil wettability. *J. Plant Nutr. Soil Sci.* 171, 14–26.
- Baisden, W. T., Amundson, R., Cook, A. C., Brenner, D. L. (2002): Turnover and storage of C and N in five density fractions from California annual grassland surface soils. *Global Biogeochem. Cycles* 16, 117–132.
- Baldock, J. A., Oades, J. M., Waters, A. G., Peng, X., Vassallo, A. M., Wilson, M. A. (1992): Aspects of the chemical structure of soil organic materials as revealed by solid-state <sup>13</sup>C NMR spectroscopy. *Biogeochem.* 16, 1–42.
- Balesdent, J. (1996): The significance of organic separates to carbon dynamics and its modelling in some cultivated soils. *Eur. J. Soil Sci.* 47, 485–493.
- Balesdent, J., Petraud, J. P., Feller, C. (1991): Effets des ultrasons sur la distribution granulométrique des matières organiques des sols. *Science du Sol* 29, 95–106.
- Balesdent, J., Besnard, E., Arrouays, D., Chenu, C. (1998): The dynamics of carbon in particle-size fractions of soil in a forest-cultivation sequence. *Plant Soil* 201, 49–57.
- Bardgett, R. D. (2005): The Biology of Soil. Oxford University Press, Oxford.
- Barois, I. (1992): Mucus production and microbial activity in the gut of two species of Amyntas (*Megascolecidae*) from cold and warm tropical climates. *Soil Biol. Biochem.* 24, 1509–1510.
- Beare, M. H., Coleman, D. C., Crossley Jr., D. A., Hendrix, P. F., Odum, E. P. (1995): A hierarchical approach to evaluating the significance of soil biodiversity to biogeochemical cycling. *Plant Soil* 170, 5–22.
- Beare, M. H., Hus, S., Coleman, D. C., Hendrix, P. F. (1997): Influences of mycelial fungi on soil aggregation and organic matter storage in conventional and no-tillage soils. *Appl. Soil Ecol.* 5, 211–219.
- Bergamaschi, B. A., Tsamakis, E., Keil, R. G., Eglinton, T. I., Montlucon, D. B., Hedges, J. I. (1997): The effect of grain size and surface area on organic matter, lignin and carbohydrate concentration, and molecular compositions in Peru Margin sediments. *Geochim. Cosmochim. Acta* 61, 1247–1260.
- Bos, R., van der Mei, H. C., Busscher, H. J. (1999): Physico-chemistry of initial microbial adhesive interactions—its mechanisms and methods for study. *FEMS Microbiol. Rev.* 23, 179–229.
- Brodowski, S., Amelung, W., Haumaier, L., Abetz, C., Zech, W. (2005): Morphological and chemical properties of black carbon in physical soil fractions as revealed by scanning electron microscopy and energy-dispersive X-ray spectroscopy. *Geoderma* 128, 116–129.
- Brodowski, S., Amelung, W., Haumaier, L., Zech, W. (2007): Black carbon contribution to stable humus in German arable soils. *Geoderma*, in press.
- Buurman, P., van Lagen, B., Piccolo, A. (2002): Increase in stability against thermal oxidation of soil humic substances as a result of self association. *Org. Geochem.* 33, 367–381.
- Chenu, C., Plante, A. F. (2006): Clay-sized organo-mineral complexes in a cultivation chronosequence: revisiting the concept of the ‘primary organo-mineral complex’. *Eur. J. Soil Sci.* 57, 596–607.
- Chenu, C., Stotzky, G. (2002): Interactions between microorganisms and soil particles: an overview, in Huang, P. M., Bollag, J.-M., Senesi, N.: Interactions between Soil Particles and Microorganisms—Impact on the Terrestrial Ecosystems. John Wiley and Sons, Chichester, UK, pp. 3–40.
- Christensen, B. T. (1992): Physical fractionation of soil and organic matter in primary particle size and density separates. *Adv. Agron.* 20, 1–90.
- Christensen, B. T. (1996): Carbon in primary and secondary organo-mineral complexes, in Carter, M. R., Stewart, B. A.: Structure and Organic Matter Storage in Agricultural Soils. Press Inc, Boca Raton, FL, USA, pp. 97–165.
- Collins, H. P., Elliott, E. T., Paustian, K., Bundy, L. G., Dick, W. A., Huggins, D. R., Smucker, A. J. M., Paul, E. A. (2000): Soil carbon pools and fluxes in long-term corn belt agroecosystems. *Soil Biol. Biochem.* 32, 157–168.
- Cuypers, C., Grotenhuis, T., Nierop, K. G. J., Franco, E. M., de Jager, A., Rulkens, W. (2002): Amorphous and condensed organic matter domains: the effect of persulfate oxidation on the composition of soil/sediment organic matter. *Chemosphere* 48, 919–931.
- Devliegher, W., Verstraete, W. (1995): *Lumbricus terrestris* in a soil core experiment—nutrient-enrichment processes (nep) and gut-associated processes (gap) and their effect on. *Soil Biol. Biochem.* 27, 1573–1580.
- Dieckow, J., Mielniczuk, J., Knicker, H., Bayer, C., Dick, D. P., Kögel-Knabner, I. (2005): Carbon and nitrogen stocks in physical fractions of a subtropical Acrisol as influenced by long-term no-till cropping systems and N fertilisation. *Plant Soil* 268, 319–328.

- Drake, H. L., Horn, M. A. (2006): Earthworms as a transient heaven for terrestrial denitrifying microbes: A review. *Eng. Life Sci.* 6, 261–265.
- Dufrene, Y. F., Boonaert, C. J. P., Rouxhet, P. G. (1996): Adhesion of *Azospirillum brasilense*: role of proteins at the cell-support interface. *Coll. Surfaces B: Biointerfaces* 7, 113–128.
- Echeverria, M. E., Markewitz, D., Morris, L. A., Hendrick, R. L. (2004): Soil Organic Matter Fractions under Managed Pine Plantations of the Southeastern USA. *Soil Sci. Soc. Am. J.* 68, 950–958.
- Egert, M., Marhan, S., Wagner, B., Scheu, S., Friedrich, M. W. (2004): Molecular profiling of 16S rRNA genes reveals diet-related differences of microbial communities in soil, gut, and casts of *Lumbricus terrestris* L. *FEMS Microbiol. Ecol.* 48, 187–197.
- Ekschmitt, K., Kandeler, E., Poll, C., Brune, A., Buscot, F., Friedrich, M., Gleixner, G., Hartmann, A., Kästner, M., Marhan, S., Miltner, A., Scheu, S., Wolters, V. (2008): Soil-carbon preservation through habitat constraints and biological limitations on decomposer activity. *J. Plant Nutr. Soil Sci.* 171, 27–35.
- Ellerbrock, R. H., Kaiser, M. (2005): Stability and composition of different soluble soil organic matter fractions—evidence from  $\delta^{13}\text{C}$  and FTIR signatures. *Geoderma* 128, 28–37.
- Ettema, C. H., Wardle, D. A. (2002): Spatial soil ecology. *Trends Ecol. Evol.* 17, 177–183.
- Eusterhues, K., Rumpel, C., Kleber, M., Kögel-Knabner, I. (2003): Stabilisation of soil organic matter by interactions with minerals as revealed by mineral dissolution and oxidative degradation. *Org. Geochem.* 34, 1591–1600.
- Eusterhues, K., Rumpel, C., Kögel-Knabner, I. (2005a): Stabilization of soil organic matter isolated via oxidative degradation. *Org. Geochem.* 36, 1567–1575.
- Eusterhues, K., Rumpel, C., Kögel-Knabner, I. (2005b): Organo-mineral associations in sandy acid forest soils: importance of specific surface area, iron oxides and micropores. *Eur. J. Soil Sci.* 56, 753–763.
- Evans, L. J., Wilson, W. G. (1985): Extractable Fe, Al and Si in B horizons of podzolic and brunisolic soils from Ontario. *Can. J. Soil Sci.* 65, 489–496.
- Flessa, H., Amelung, W., Helfrich, M., Wiesenberg, G. L. B., Gleixner, G., Brodowski, S., Rethemeyer, J., Kramer, C., Grootes, P.-M. (2008): Storage and stability of organic matter and fossil carbon in a Luvisol and Phaeozem with continuous maize cropping: A synthesis. *J. Plant Nutr. Soil Sci.* 171, 36–51.
- Fry, S. (2007): Spatial distribution of soil organisms, in Paul, E. A. (ed.): *Soil Microbiology, Biochemistry and Soil Ecology*. 3rd edn., Elsevier, San Diego, in press.
- Gaillard, V., Chenu, C., Recous, S. (2003): Carbon mineralisation in soil adjacent to plant residues of contrasting biochemical quality. *Soil Biol. Biochem.* 35, 93–99.
- Gaston, L. A., Locke, M. A. (2002): Differences in microbial biomass, organic carbon, and dye sorption between flow and no-flow regions of unsaturated soil. *J. Environ. Qual.* 31, 1406–1408.
- Gleixner, G., Danier, H.-J., Werner, R. A., Schmidt, H. L. (1993): Correlations between the  $^{13}\text{C}$  content of primary and secondary plant-products in different cell compartments and that in decomposing basidiomycetes. *Plant Physiol.* 102, 1287–1290.
- Golchin, A., Oades, J. M., Skjemstad, J. O., Clake, P. (1994): Study of free and occluded particulate organic matter in soils by solid-state  $^{13}\text{C}$  CP/MAS NMR spectroscopy and scanning electron microscopy. *Aust. J. Soil. Res.* 32, 285–309.
- Griffith, S. M., Schnitzer, M. (1977): Organic compounds formed by the hydrogen peroxide oxidation of soils. *Can. J. Soil. Sci.* 57, 223–231.
- Guggenberger, G., Christensen, B. T., Zech, W. (1994): Land-use effects on the composition of organic matter in particle-size separates of soils: I. Lignin and carbohydrate signature. *Eur. J. Soil Sci.* 45, 449–458.
- Hassink, J. (1997): The capacity of soils to preserve organic C and N by their association with clay and silt particles. *Plant Soil* 191, 77–87.
- Hedges, J. I., Oades, J. M. (1997): Comparative organic geochemistries of soils and marine sediments. *Org. Geochem.* 27, 319–361.
- Helfrich, M., Flessa, H., Mikutta, R., Dreves, A., Ludwig, B. (2007): Assessing the suitability of chemical fractionation methods for extracting stable soil organic matter pools using  $^{13}\text{C}$  and  $^{14}\text{C}$  analysis, *Eur. J. Soil Sci.*, submitted.
- Hindell, R. P., McKenzie, B. M., Tisdall, J. M. (1997): Destabilization of soil during the production of earthworm (Lumbricidae) and artificial casts. *Biol. Fertil. Soils* 24, 153–163.
- Horn, M. A., Mertel, R., Gehre, M., Kaestner, M., Drake, H. L. (2006): In vivo emission of dinitrogen by earthworms via denitrifying bacteria in the gut. *Appl. Environ. Microbiol.* 72, 1013–1018.
- Huang, Y., Li, B. C., Bryant, C., Bol, R., Eglinton, G. (1999): Radiocarbon dating of aliphatic hydrocarbons: A new approach for dating passive-fraction carbon in soil horizons. *Soil Sci. Soc. Am. J.* 63, 1181–1187.
- Janzen, H. H., Campbell, C. A., Brandt, S. A., Lafond, G. P., Toenley-Smith, L. (1992): Light fraction organic matter in soils from long term crop rotations. *Soil Sci. Soc. Am. J.* 56, 1799–1806.
- Jocteur Monrozier, L., Ladd, J. N., Fitzpatrick, R. W., Foster, R. C., Raupach, M. (1991): Components and microbial biomass content of size fractions in soils of contrasting aggregation. *Geoderma* 49, 37–62.
- John, B., Yamashita, T., Ludwig, B., Flessa, H. (2005): Storage of organic carbon in aggregate and density fractions of silty soils under different types of land use. *Geoderma* 128, 63–79.
- Kahle, M., Kleber, M., Jahn, R. (2002): Carbon storage in loess derived surface soils from Central Germany: Influence of mineral phase variables. *J. Plant Nutr. Soil Sci.* 165, 141–149.
- Kahle, M., Kleber, M., Jahn, R. (2003a): Retention of dissolved organic matter by illitic soils and clay fractions: Influence of mineral phase properties. *J. Plant Nutr. Soil Sci.* 166, 737–741.
- Kahle, M., Kleber, M., Torn, M. S., Jahn, R. (2003b): Carbon storage in coarse and fine clay fractions of illitic soils. *Soil Sci. Soc. Am. J.* 67, 1732–1739.
- Kaiser, K., Guggenberger, G. (2003): Mineral surfaces and soil organic matter. *Eur. J. Soil Sci.* 54, 219–236.
- Kaiser, K., Guggenberger, G. (2007a): Sorptive stabilization of organic matter by microporous goethite: sorption into small pores vs. surface complexation. *Eur. J. Soil Sci.* 53, 639–644, doi: 10.1111/j.1365-2389.2006.00799.x.
- Kaiser, K., Guggenberger, G. (2007b): Distribution of hydrous aluminium and iron over density fractions depends on organic matter load and ultrasonic dispersion. *Geoderma*, in press.
- Kaiser, K., Zech, W. (1996): Defects in estimation of aluminum in humus complexes of podzolic soils by pyrophosphate extraction. *Soil Sci.* 161, 452–458.
- Kaiser, K., Mikutta, R., Guggenberger, G. (2007): Increased stability of organic matter sorbed to ferrihydrite and goethite upon aging. *Soil Sci. Soc. Am. J.*, in press.
- Kalbitz, K., Schmerwitz, J., Schwesig, D., Matzner, E. (2003): Biodegradation of soil-derived dissolved organic matter as related to its properties. *Geoderma* 113, 273–291.

- Kalbitz, K., Schwesig, D., Rethemeyer, J., Matzner, E. (2005): Stabilization of dissolved organic matter by sorption to the mineral soil. *Soil Biol. Biochem.* 37, 1319–1331.
- Kanazawa, S., Filip, Z. (1986) Distribution of microorganisms, total biomass and enzyme activities in different particles of brown soil. *Microb. Ecol.* 12, 205–215.
- Kandeler, E., Dick, R. P. (2006): Distribution and Function of Soil Enzymes in Agroecosystems, in Benckiser, G., Schnell, S.: Biodiversity in Agricultural Production Systems. Elsevier, Amsterdam, in press.
- Kandeler, E., Stemmer, M., Klimanek, E. M. (1999a): Response of soil microbial biomass, urease and xylanase within particle size fractions to long-term soil management. *Soil Biol. Biochem.* 31, 261–273.
- Kandeler, E., Palli, S., Stemmer, M., Gerzabek, M. H. (1999b): Tillage changes microbial biomass and enzyme activities in particle-size fractions of a Haplic Chernozem. *Soil Biol. Biochem.* 31, 1253–1264.
- Kandeler, E., Tschirko, D., Spiegel, H. (1999c): Long-term monitoring of microbial biomass, N-mineralization and enzyme activities of a Chernozem under different tillage management. *Biol. Fertil. Soils* 28, 343–351.
- Kandeler, E., Tschirko, D., Bruce, K. D., Stemmer, M., Hobbs, P. J., Bardgett, R. D., Amelung, W. (2000): The structure and function of the soil microbial community in microhabitats of a heavy metal polluted soil. *Biol. Fertil. Soils* 32, 390–400.
- Kandeler, E., Tschirko, D., Stemmer, M., Schwarz, S., Gerzabek, M. H. (2001): Organic matter and soil microorganisms – Investigations from the micro- to the macro-scale. *Die Bodenkultur* 52, 117–131.
- Kandeler, E., Stemmer, M., Gerzabek, M. H. (2005): Role of microorganisms in carbon cycling in soils, in Buscot, F., Varma, A.: Microorganisms in soils: roles in genesis and functions. Springer-Verlag, Berlin, Heidelberg, pp. 139–157.
- Kassim, J. K., Gafoor, S. N., Adams, W. A. (1984): Ferrihydrite in pyrophosphate extracts of Podzol B horizons. *Clay Min.* 19, 99–106.
- Keil, R. G., Montlucon, D. B., Prah, F. G., Hedges, J. I. (1994): Sorptive preservation of labile organic matter in marine sediments. *Nature* 370, 549–552.
- Kiem, R., Kögel-Knabner, I. (2003): Contribution of lignin and polysaccharides to the refractory carbon pool as studied in C-depleted arable soils. *Soil Biol. Biochem.* 35, 101–118.
- Kirchmann, H., Haberhauer, G., Kandeler, E., Sessitsch, A., Gerzabek, M. H. (2004): Level and quality of organic matter input regulates biogeochemistry of soils – Synthesis of a long-term agricultural field study. *Global Biogeochem. Cycles* 18, GB4011, doi:10.1029/2003GB002204.
- Kleber, M., Mertz, C., Zikeli, S., Knicker, H., Jahn, R. (2004): Changes in surface reactivity and organic matter composition of clay subfractions with duration of fertilizer deprivation. *Eur. J. Soil Sci.* 55, 381–391.
- Kleber, M., Mikutta, R., Torn, M. S., Jahn, R. (2005): Poorly crystalline mineral phases protect organic matter in acid subsoil horizons. *Eur. J. Soil Sci.* 56, 717–725.
- Kleber, M., Sollins, P., Sutton, R. (2007): A conceptual model of organo-mineral interactions in soils: self-assembly of organic molecular fragments into multilayered structures on mineral surfaces. *Biogeochem.*, accepted.
- Knicker, H. (2004): Stabilization of N-compounds in soil and organic-matter-rich sediments—what is the difference? *Mar. Chem.* 92, 167–195.
- Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E., Marschner, B., von Lützow, M. (2008): An integrative approach of organic matter stabilization in temperate soils: Linking chemistry, physics, and biology. *J. Plant Nutr. Soil Sci.* 171, 5–13.
- Kubienna, W. L. (1948): Entwicklungslehre des Bodens. Springer, Wien.
- Lavelle, P., Pashanasi, B., Charpentier, F., Gilot, C., Rossi, J.-P., Derouard, L., Andre, J., Ponge, J.-F., Bernier, N. (2004): Effects of earthworms on soil organic matter dynamics at a landscape scale over decades, in: Edwards C. A. (ed.), *Earthworm Ecology*. 2nd edn., CRC Press, Boca Raton, pp. 145–160.
- Leifeld, J., Kögel-Knabner, I. (2001): Organic carbon and nitrogen in fine soil fractions after treatment with hydrogen peroxide. *Soil Biol. Biochem.* 33, 2155–2158.
- Mahieu, N., Powelson, D. S., Randall, E. W. (1999): Statistical analysis of published carbon-13 CPMAS NMR spectra of soil organic matter. *Soil Sci. Soc. Am. J.* 63, 307–319.
- Marhan, M., Scheu, S. (2006): Mixing of different mineral soil layers by endogeic earthworms affects carbon and nitrogen mineralization. *Biol. Fertil. Soils* 32, 308–314.
- Marhan, S., Kandeler, E., Scheu, S. (2007): Phospholipid fatty acid profiles and xylanase activity in particle size fractions of forest soil and casts of *Lumbricus terrestris* L. (Oligochaeta, Lumbricidae). *Appl. Soil Ecol.* 35, 412–422.
- Marschner, B., Brodowski, S., Dreves, A., Gleixner, G., Gude, A., Grootes, P.-M., Hamer, U., Heim, A., Jandl, G., Ji, R., Kaiser, K., Kalbitz, K., Kramer, C., Leinweber, P., Rethemeyer, J., Schäffer, A., Schmidt, M. W. I., Schwark, L., Wiesenberger, G. L. B. (2008): How relevant is recalcitrance for the stabilization of organic matter in soils? *J. Plant Nutr. Soil Sci.* 171, 91–110.
- Martin, A. (1991): Short-term and long-term effects of the endogenic earthworm *Millsonia anomala* (Omodeo) (Megascolecidae, Oligochaeta) of tropical savannas on soil organic matter. *Biol. Fertil. Soils* 11, 234–238.
- Martin, F., Saint-Jimenez, C., Gonzalez-Vila, F. J. (1981): The persulfate oxidation of a soil humic acid. *Soil Sci.* 132, 200–203.
- Marx, M. C., Kandeler, E., Wood, M., Wermbter, N., Jarvis, S. C. (2005): Exploring the enzymatic landscape: distribution and kinetics of hydrolytic enzymes in soil particle size fractions. *Soil Biol. Biochem.* 37, 35–48.
- Masiello, C. A., Chadwick, O. A., Southon, J., Torn, M. S., Harden, J. W. (2004): Weathering controls on mechanisms of carbon storage in grassland soils. *Global Biogeochem. Cycles* 18, GB4023, doi:10.1029/2004GB002219.
- Mayer, L. M. (1994a): Relationships between mineral surfaces and organic carbon concentrations in soils and sediments. *Chem. Geol.* 114, 347–363.
- Mayer, L. M. (1994b): Surface area control of organic carbon accumulation in continental shelf sediments. *Geochim. Cosmochim. Acta* 58, 1271–1284.
- Mayer, L. M., Xing, B. (2001): Organic matter–surface area relationships in acid soils. *Soil Sci. Soc. Am. J.* 65, 250–258
- Mayer, L. M., Schick, L. L., Hardy, K. R., Wagai, R., McCarthy, J. (2004): Organic matter in small mesopores in sediments and soils. *Geochim. Cosmochim. Acta* 68, 3863–3872.
- Mikutta, C., Lang, F., Kaupenjohann, M. (2004): Soil organic matter clogs mineral pores: Evidence from H-1-NMR and N-2 adsorption. *Soil Sci. Soc. Am. J.* 68, 1853–1862.
- Mikutta, R., Kleber, M., Kaiser, K., Jahn, R. (2005): Review: organic matter removal from soils using hydrogen peroxide, sodium hypochlorite, and disodium peroxodisulfate. *Soil Sci. Soc. Am. J.* 69, 120–135.

- Mikutta, R., Kleber, M., Torn, M. S., Jahn, R. (2006): Stabilization of soil organic matter: Association with minerals or chemical recalcitrance? *Biogeochem.* 77, 25–56.
- Mikutta, R., Mikutta, C., Kalbitz, K., Scheel, T., Kaiser, K., Jahn, R. (2007): Biodegradation of forest floor organic matter bound to minerals via different binding mechanisms. *Geochim. Cosmochim. Acta* 71, 2569–2590.
- Mummey, D. L., Rillig, M. C., Six, J. (2006): Endogeic earthworms differentially influence bacterial communities associated with different soil aggregate size fractions. *Soil Biol. Biochem.* 38, 1608–1614.
- Murphy, E. M., Zachara, J. M., Smith, S. C. (1990): Influence of mineral-bound humic substances on the sorption of hydrophobic organic compounds. *Environ. Sci. Technol.* 24, 1507–1516.
- North, P. F. (1976): Towards an absolute measurement of soil structural stability using ultrasound. *J. Soil Sci.* 27, 451–459.
- Nunan, N., Wu, K. J., Young, I. M., Crawford, J. W., Ritz, K. (2003): Spatial distribution of bacterial communities and their relationships with the micro-architecture of soil. *FEMS Microbiol. Ecol.* 44, 203–215.
- Oades, J. M. (1988): The retention of organic matter in soils. *Biogeochem.* 5, 35–70.
- Omoike, A., Chorover, J. (2006): Adsorption to goethite of extracellular polymeric substances from *Bacillus subtilis*. *Geochim. Cosmochim. Acta* 70, 827–838.
- Paul, E. A., Follett, R. F., Leavitt, S. W., Halvorson, A., Peterson, G. A., Lyon, D. J. (1997): Radiocarbon dating for determination of soil organic matter pool sizes and dynamics. *Soil Sci. Soc. Am. J.* 61, 1058–1067.
- Paul, E. A., Collins, H. P., Leavitt, S. W. (2001): Dynamics of resistant soil carbon of midwestern agricultural soils measured by naturally occurring C-14 abundance. *Geoderma* 104, 239–256.
- Paul, E. A., Morris, S. J., Conant, R. T., Plante, A. F. (2006): Does the acid hydrolysis-incubation method measure meaningful soil organic carbon pools? *Soil Sci. Soc. Am. J.* 70, 1023–1035.
- Plante, A. F., Conant, R. T., Paul, E. A., Paustian, K., Six, J. (2006): Acid hydrolysis of easily dispersed and microaggregate-derived silt- and clay-sized fractions to isolate resistant soil organic matter. *Eur. J. Soil Sci.* 57, 456–467.
- Poll, C., Thiede, A., Wermbter, N., Sessitsch, A., Kandeler, E. (2003): Micro-scale distribution of microorganisms and microbial enzyme activities in a soil with long-term organic amendment. *Eur. J. Soil Sci.* 54, 715–724.
- Poll, C., Ingwersen, J., Stemmer, M., Gerzabek, M. H., Kandeler, E. (2006): Mechanisms of solute transport affect small-scale abundance and function of soil microorganisms in the detritusphere. *Eur. J. Soil Sci.* 57, 583–595.
- Poll, C., Marhan, S., Ingwersen, J., Kandeler, E. (2007): Dynamics of carbon turnover and microbial abundance in a rye detritusphere. *Soil Biol. Biochem.*, in press.
- Rasmussen, C., Torn, M. S., Southard, R. J. (2005): Mineral assemblage and aggregates control carbon dynamics in a California conifer forest. *Soil Sci. Soc. Am. J.* 69, 1711–1721.
- Roscoe, R., Buurman, P. (2003): Tillage effects on soil organic matter in density fractions of a Cerrado Oxisol. *Soil Tillage Res.* 70, 107–119.
- Rovira, P., Vallejo, V. R. (2003): Physical protection and biochemical quality of organic matter in mediterranean calcareous forest soils: a density fractionation approach. *Soil Biol. Biochem.* 35, 245–261.
- Rumpel, C., Eusterhues, K., Kögel-Knabner, I. (2004): Location and chemical composition of stabilized organic carbon in topsoil and subsoil horizons of two acid forest soils. *Soil Biol. Biochem.* 36, 177–190.
- Saggar, S., Parshotam, A., Sparling, G. P., Feltham, C. W., Hart, P. B. S. (1996): <sup>14</sup>C-labelled ryegrass turnover and residence times in soils varying in clay content and mineralogy. *Soil Biol. Biochem.* 28, 1677–1686.
- Sampedro, L., Jeannotte, R., Whalen, J. K. (2006): Trophic transfer of fatty acids from gut microbiota to the earthworm *Lumbricus terrestris* L. *Soil Biol. Biochem.* 38, 2188–2198.
- Scheel, T., Dörfler, C., Kalbitz, K. (2007): Precipitation of dissolved organic matter by Al stabilizes carbon in acidic forest soils. *Soil Sci. Soc. Am. J.* 71, 64–74.
- Scheu, S. (1987): The role of substrate feeding earthworms (Lumbricidae) for bioturbation in a beechwood soil. *Oecologia* 72, 192–196.
- Schmid, M., Selesi, D., Rothballer, M., Schloter, M., Natuschka, L., Kandeler, E., Hartmann, A. (2006): Localization and visualisation of microbial population structure and activity in soil microhabitats, in König/Varma: Intestinal Microorganisms of Termites and other Invertebrates. Soil Biology-Serie, Vol. 6, Springer Verlag, Berlin Heidelberg, pp. 439–462.
- Schmidt, M. W. I., Rumpel, C., Kögel-Knabner, I. (1999a): Evaluation of an ultrasonic dispersion method to isolate primary organo-mineral complexes from soils. *Eur. J. Soil Sci.* 50, 87–94.
- Schmidt, M. W. I., Rumpel, C., Kögel-Knabner, I. (1999b): Particle size fractionation of soil containing coal and combusted particles. *Eur. J. Soil Sci.* 50, 515–522.
- Schmidt, M. W. I., Skjemstad, J. O., Gehrt, E., Kögel-Knabner, I. (1999c): Charred organic carbon in German chernozemic soils. *Eur. J. Soil Sci.* 50, 351–365.
- Schmidt, O., Scrimgeour, C. M., Curry, J. P. (1999d): Carbon and nitrogen stable isotope ratios in body tissue and mucus of feeding and fasting earthworms (*Lumbricus festivus*). *Oecologia* 118, 9–15.
- Schönholzer, F., Hahn, D., Zeyer, J. (1999): Origins and fate of fungi and bacteria in the gut of *Lumbricus terrestris* L. studied by image analysis. *FEMS Microbiol. Ecol.* 28, 235–248.
- Schöning, I., Kögel-Knabner, I. (2006): Chemical composition of young and old carbon pools throughout Cambisol and Luvisol profiles under forest. *Soil Biol. Biochem.* 38, 2411–2424.
- Schöning, I., Morgenroth, G., Kögel-Knabner, I. (2005a): O/N-alkyl and alkyl C are stabilised in fine particle size fractions of forest soils. *Biogeochem.* 73, 475–497.
- Schöning, I., Knicker, H., Kögel-Knabner, I. (2005b): Intimate associations between O/N-alkyl carbon and iron oxides in clay fractions of forest soils. *Org. Geochem.* 36, 1378–1390.
- Schulten, H.-R., Leinweber, P. (1999): Thermal stability and composition of mineral-bound organic matter in density fractions of soil. *Eur. J. Soil Sci.* 50, 237–248.
- Schulten, H.-R., Leinweber, P. (2000): New insights in organo-mineral particles: composition, properties and models of molecular structure. *Biol. Fertil. Soils* 30, 399–432.
- Schulten, H.-R., Leinweber, P., Theng, B. K. G. (1996): Characterization of organic matter in an interlayer-organic complex from soil by pyrolysis methylation-mass spectrometry. *Geoderma* 69, 105–118.
- Schuppli, P. A., Ross, G. J., McKeague, J. A. (1983): The effective removal of suspended materials from pyrophosphate extracts of soils from tropical and temperate regions. *Soil Sci. Soc. Am. J.* 47, 1026–1032.

- Scullion, J., Malik, A., (2000): Earthworm activity affecting organic matter, aggregation and microbial activity in soils restored after opencast mining for coal. *Soil Biol. Biochem.* 32, 119–126.
- Selesi, D., Pattis, I., Schmid, M., Kandeler, E., Hartmann, A. (2007): Quantification of bacterial RubisCO genes in soil by cbbL targeted real-time PCR. *J. Microbiol. Methods*, 69, 497–503, doi:10.1016/j.mimet.2007.03.002.
- Sessitsch, A., Weilharter, A., Gerzabek, M. H., Kirchmann, H., Kandeler, E. (2001): Microbial population structures in soil particle size fractions of a long-term fertilizer field experiment. *Appl. Environ. Microbiol.* 67, 4215–4224.
- Shaw, C., Pawluk, S. (1986): The development of soil structure by *Octolasion tyraeum*, *Aporrectodea turgida* and *Lumbricus terrestris* in parent materials belonging to different textural classes. *Pedobiologia* 29, 327–339.
- Shipitalo, M. J., Protz, R. (1988): Factors influencing the dispersibility of clay in worm casts. *Soil Sci. Soc. Am. J.* 52, 764–769.
- Shipitalo, M. J., Protz, R. (1989): Chemistry and micromorphology of aggregation in earthworm casts. *Geoderma* 45, 357–374.
- Six, J., Jastrow, J. D. (2002): Organic matter turnover, in Lal, R.: Encyclopedia of Soil Science. Marcel Dekker, New York, pp. 936–942.
- Six, J., Elliott, E. T., Paustian, K., Doran, J. W. (1998): Aggregation and soil organic matter accumulation in cultivated and native grassland soils. *Soil Sci. Soc. Am. J.* 62, 1367–1377.
- Six, J., Conant, R. T., Paul, E. A., Paustian, K. (2002): Stabilization mechanisms of soil organic matter: Implications for C-saturation of soils. *Plant Soil* 241, 155–176.
- Skjemstad, J. O., Le Feuvre, R. P., Prebble, R. E. (1990): Turnover of soil organic matter under pasture as determined by <sup>13</sup>C natural abundance. *Aust. J. Soil Res.* 28, 267–276.
- Sollins, P., Swanston, C., Kleber, M., Filley, T., Kramer, M., Crow, S., Caldwell, B. A., Lajtha, K., Bowden, R. (2006): Organic C and N stabilization in a forest soil: Evidence from sequential density fractionation. *Soil Biol. Biochem.* 38, 3313–3324.
- Speir, T. W., Ross, D. J. (2002): Hydrolytic enzyme activities to assess soil degradation and recover, in Burns, R. G., Dick, R. P.: Enzymes in the Environment: Activity, Ecology and Applications. Marcel Dekker, New York, pp. 407–431.
- Stemmer, M., Gerzabek, M. H., Kandeler, E. (1998): Organic matter and enzyme activity in particle size fractions of soils obtained after low-energy sonication. *Soil Biol. Biochem.* 30, 9–17.
- Stemmer, M., Gerzabek, M. H., Kandeler, E. (1999): Invertase and xylanase activity of bulk soil and particle-size fractions during maize straw decomposition. *Soil Biol. Biochem.* 31, 9–18.
- Stevenson, F. J. (1982): Humus Chemistry. Wiley, New York.
- Swanston, C. W., Tron, M. S., Hanson, P. J., Southon, J. R., Garten, C. T., Hanlon, E. M., Ganio, L. (2005): Initial characterization of processes of soil carbon stabilization using forest stand-level radiocarbon enrichment. *Geoderma* 128, 52–62.
- Theng, B. K. G. (1979): Formation and properties of clay-polymer complexes. Elsevier Science Publishing Co., New York.
- Theng, B. K. G., Churchman, G. J., Newman, R. H. (1986): The occurrence of interlayer clay-organic complexes in two New Zealand soils. *Soil Sci.* 142, 262–266.
- Tiessen, H., Stewart, J. W. B., Hunt, H. W. (1984): Concepts of soil organic matter transformations in relation to organo-mineral particle-size fractions. *Plant Soil* 76, 287–295.
- Tjunov, A. V., Scheu, S. (2000): Microbial biomass, biovolume and respiration in *Lumbricus terrestris* L. cast material of different age. *Soil Biol. Biochem.* 32, 265–275.
- Torn, M. S., Trumbore, S. E., Chadwick, O. A., Vitousek, P. M., Hendricks, D. M. (1997): Mineral control of soil organic carbon storage and turnover. *Nature* 389, 170–173.
- Turchenek, L. W., Oades, J. M. (1979): Fractionation of organo-mineral complexes by sedimentation and density techniques. *Geoderma* 21, 311–343.
- Vanyushina, A. Y., Travnikova, L. S. (2003): Organic-mineral interactions in soils: a review. *Euras. Soil Sci.* 36, 379–387.
- von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., Flessa, H. (2006): Stabilization of organic matter in temperate soils: Mechanisms and their relevance under different soil conditions—a review. *Eur. J. Soil Sci.* 57, 426–445.
- von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E., Marschner, B. (2007): SOM fractionation methods: Relevance to functional pools and to stabilization mechanisms—a review. *Soil Biol. Biochem.*, in press.
- Wander, M. (2004): Soil organic matter fractions and their relevance to soil function, in Weil, R. R.: Soil organic matter in sustainable agriculture. CRC Press, Boca Raton, FL, USA, pp. 67–102.
- Watt, M., Silk, W. K., Passioura, J. B. (2006): Rates of root and organism growth, soil conditions, and temporal and spatial development of the rhizosphere. *Ann. Bot.* 97, 839–855.
- Wattel-Koekkoek, E. J. W., Buurman, P. (2004): Mean residence time of kaolinite and smectite-bound organic matter in Mozambiquan soils. *Soil Sci. Soc. Am. J.* 68, 154–161.
- Weidler, P. G., Degovics, G., Laggner, P. (1998): Surface roughness created by acidic dissolution of synthetic goethite monitored by SAXS and N<sub>2</sub>-adsorption isotherms. *J. Col. Interf. Sci.* 197, 1–8.
- Zimmerman, A. R., Goyne, K. W., Chorover, J., Komarneni, S., Brantley, S. L. (2004): Mineral mesopore effects on nitrogenous organic matter adsorption. *Org. Geochem.* 35, 355–375.